

Spectroscopy of tetraquark states.

Elena Santopinto* and Giuseppe Galatà

INFN and Università di Genova, via Dodecaneso 33, 16142 Genova, Italy

A complete classification of $qq\bar{q}\bar{q}$ tetraquark states in terms of the spin-flavor, color and spatial degrees of freedom has been constructed. The permutation symmetry properties of both the spin-flavor and orbital parts of the qq and $\bar{q}\bar{q}$ subsystems are discussed. This complete classification is general and model independent and it is useful both for model builders and experimentalists. The total wave functions are also explicitly constructed in the hypothesis of ideal mixing; this basis for tetraquark states will enable the eigenvalue problem to be solved for a definite dynamical model. An evaluation of the tetraquark spectrum is obtained from the Iachello mass formula for normal mesons, here generalized to tetraquark systems. This mass formula is a generalization of the Gell-Mann Okubo mass formula, whose coefficients have been upgraded by a study of the latest PDG data. The ground state tetraquark nonet is identified with $f_0(600), \kappa(800), f_0(980), a_0(980)$. The diquark-antidiquark limit is also studied.

PACS numbers: 14.40.Cs, 12.39.-x, 02.20.-a

I. INTRODUCTION

The KLOE, E791 and BES collaborations have recently provided evidence of the low mass resonances $f_0(600)$ [1, 2, 3], formerly called $\sigma(450)$, and $\kappa(800)$ [2, 3], triggering new interest in meson spectroscopy. Maiani et al. [4] have suggested that the lowest lying scalar mesons, $f_0(980)$, $a_0(980)$, $\kappa(800)$ and $f_0(600)$ could be described not as $q\bar{q}$ states, but as more complex tetraquark states, in particular as two clusters of two quarks and two antiquarks, i.e. a diquark and antidiquark system. The quark-antiquark assignment to P-waves [5] has never really worked in the scalar case [6, 7]. Moreover, the $f_0(980)$ is more associated to strange than to up or down quarks as can be inferred from its higher mass and its decays [4, 6, 7], while in a simple quark-antiquark scheme it is associated with non-strange quarks [5]; for this reason it is difficult to explain both its mass and its decay properties [4, 6, 7] at the same time. One of the arguments by Jaffe [6, 7] and Maiani [4] against the hypothesis of simple $q\bar{q}$ states is the observation that the experimental mass spectrum corresponding to this nonet is like a parabola with a maximum in the centre of the nonet corresponding to the $f_0(980)$ and $a_0(980)$, while in the $q\bar{q}$ case the parabola would be reversed and so the maximum would be at the edge of the nonet.

Other identifications have been proposed [8], in particular quasi molecular-states (see Refs. [9, 10, 11] and references therein, [5]) and uncorrelated $qq\bar{q}\bar{q}$ [12, 13]. Previous works on heavy tetraquark mesons can be found in Refs. [14, 15, 16, 17] and for light mesons in [18, 19, 20] and references therein. As early as the 1970s Jaffe studied tetraquark systems in a bag model and discussed the resulting rich spectrum together with the problem of the missing resonances [6, 7, 12]. For review articles both on the experiments and on the theoretical models we refer the reader to [8, 21, 22].

In this article, we address the problem of constructing a complete classification scheme of the two quark-two antiquark states in terms of $SU_{sf}(6)$. We identify the representations that contain exotics, i.e. states that cannot be constructed by $q\bar{q}$ only. The tetraquark $O(3) \otimes SU_{sf}(6) \otimes SU_c(3)$ wave functions are explicitly constructed for the first time. They should be color singlets and, since they are composed of two quarks and two antiquarks, i. e. two couples of identical fermions, they should be antisymmetric for the exchange of the two quarks and the two antiquarks. The permutation symmetry properties of both the spin-flavor and the orbital parts of the qq and $\bar{q}\bar{q}$ subsystems are discussed. The total wave functions are also explicitly constructed in the ideal mixing hypothesis, and can be useful in order to construct tetraquark models. Finally, an evaluation of the tetraquark spectrum for the lowest scalar mesons is obtained from a generalization of the Iachello mass formula for normal mesons [23].

The classification of the states is general and is valid whichever dynamical model for tetraquarks is chosen. As an application, in section V we develop a simple diquark-antidiquark model with no spatial excitations inside diquarks. The states are a subset of the general case.

*Electronic address: elena.santopinto@ge.infn.it

II. THE CLASSIFICATION OF TETRAQUARK STATES

As for all multiquark systems, the tetraquark wave function contains contributions connected to the spatial degrees of freedom and the internal degrees of freedom of color, flavor and spin. In order to classify the corresponding states, we shall make use as much as possible of symmetry principles without, for the moment, introducing any explicit dynamical model. In the construction of the classification scheme we are guided by two conditions: the tetraquark wave functions should be a color singlet, as all physical states, and since tetraquarks are composed of two couples of identical fermions, their states must be antisymmetric for the exchange of the two quarks and the two antiquarks.

In the following, we adopt the usual notation $[R]$ for the representations, where R is the dimension of the representation.

A. The $SU(3)_f$ -flavor classification of $qq\bar{q}\bar{q}$ states

The allowed $SU(3)_f$ representations for the $qq\bar{q}\bar{q}$ mesons are obtained by means of the product

$$[3] \otimes [\bar{3}] \otimes [3] \otimes [\bar{3}] = [1] \oplus [8] \oplus [8] \oplus [10] \oplus [8] \oplus [8] \oplus [1] \oplus [\bar{10}] \oplus [27] \quad (1)$$

The allowed isospin values are $I = 0, \frac{1}{2}, 1, \frac{3}{2}, 2$, while the hypercharge values are $Y = 0, \pm 1, \pm 2$. We can notice that the values $I = \frac{3}{2}, 2$ and $Y = \pm 2$ are exotic, which means that they are forbidden for the $q\bar{q}$ mesons.

In Appendix A the flavor states in the $qq\bar{q}\bar{q}$ configuration are explicitly written.

B. The $SU(3)_c$ -color classification of $qq\bar{q}\bar{q}$ states

Color representations for $qq\bar{q}\bar{q}$ mesons are those written in (1) for the flavor case. However, the only color representation allowed for mesons (or in general for any isolated particle) is the singlet, so there are two colour representations for $qq\bar{q}\bar{q}$ mesons, while there is only one singlet for normal mesons. This fact implies that color for tetraquarks is not a trivial quantum number as it was for conventional mesons.

C. The $SU(2)_s$ -spin classification of $qq\bar{q}\bar{q}$ states

The $qq\bar{q}\bar{q}$ spin states are given by the product

$$[2] \otimes [2] \otimes [2] \otimes [2] = [1] \oplus [3] \oplus [1] \oplus [3] \oplus [3] \oplus [5] \quad (2)$$

We can see that tetraquarks can have an exotic spin $S = 2$, value forbidden for $q\bar{q}$ mesons.

In Appendix B the spin states in the $qq\bar{q}\bar{q}$ configuration are explicitly written.

D. The $SU(6)_{sf}$ -spin-flavor classification of $qq\bar{q}\bar{q}$ states

The spin-flavor $SU(6)_{sf}$ $qq\bar{q}\bar{q}$ representations are obtained by means of the product

$$[6] \otimes [\bar{6}] \otimes [6] \otimes [\bar{6}] = [1] \oplus [35] \oplus [35] \oplus [405] \oplus [35] \oplus [280] \oplus [35] \oplus [1] \oplus [\bar{280}] \oplus [189] \quad (3)$$

A complete classification of the tetraquark states involves the analysis of the flavor and spin content of each spin-flavor representation, i.e. the decomposition of the representation of $SU(6)_{sf}$ into those of $SU(3)_f \otimes SU(2)_s$ in the notation [flavor repr., spin repr.],

$$[189] = [8, 5] \oplus [10, 3] \oplus [27, 1] \oplus [\bar{10}, 3] \oplus 2[8, 3] \oplus [8, 1] \oplus [1, 1] \oplus [1, 5] \quad (4)$$


$$[280] = [10, 5] \oplus [8, 5] \oplus [27, 3] \oplus [10, 3] \oplus 2[8, 3] \oplus [10, 1] \oplus [\bar{10}, 1] \oplus [8, 1] \oplus [1, 3] \quad (5)$$

$$[\bar{280}] = [\bar{10}, 5] \oplus [8, 5] \oplus [27, 3] \oplus [\bar{10}, 3] \oplus 2[8, 3] \oplus [\bar{10}, 1] \oplus [10, 1] \oplus [8, 1] \oplus [1, 3] \quad (6)$$

$$[405] = [1, 1] \oplus [1, 5] \oplus [8, 5] \oplus 2[8, 3] \oplus [27, 1] \oplus [8, 1] \oplus [27, 3] \oplus [10, 3] \oplus [\bar{10}, 3] \oplus [27, 5] \quad (7)$$

$$[1] = [1, 1] \quad (8)$$

$$[35] = [1, 3] \oplus [8, 1] \oplus [8, 3] \quad (9)$$

Figure 1: $qq\bar{q}\bar{q}$ mesons' total angular momentum scheme

E. Angular momentum, parity and charge conjugation quantum numbers

The total angular momentum, parity and charge conjugation quantum numbers for the $q\bar{q}$ mesons are well known. Thus, here we recall only that the following J^{PC} combinations are forbidden for normal mesons:

$$0^{--}, (even)^{+-}, (odd)^{-+}. \quad (10)$$

Tetraquarks are made up of four objects, so we have to define three relative coordinates (see figure 1) [24]

$$\vec{r}_{13} = \vec{r}_3 - \vec{r}_1 \quad (11a)$$

$$\vec{r}_{24} = \vec{r}_4 - \vec{r}_2 \quad (11b)$$

$$\vec{r}_{12-34} = \vec{r}_{CM24} - \vec{r}_{CM13} = \frac{m_2\vec{r}_2 + m_4\vec{r}_4}{m_2 + m_4} - \frac{m_1\vec{r}_1 + m_3\vec{r}_3}{m_1 + m_3}. \quad (11c)$$

This is only a possible choice of coordinates. Other types of coordinates, useful to describe the strong decays, can be defined [18].

In the tetraquark case, we have four different spins and three orbital angular momenta. The total angular momentum J can be obtained by combining spins and orbital momenta, as shown in figure 1.

The parity for a tetraquark system is the product of the intrinsic parities of the quarks and the antiquarks times the factors coming from the spherical harmonics [24].

$$P = P_q P_q P_{\bar{q}} P_{\bar{q}} (-1)^{L_{13}} (-1)^{L_{24}} (-1)^{L_{12-34}} = (-1)^{L_{13} + L_{24} + L_{12-34}} \quad (12)$$

Using our coordinates, tetraquark charge conjugation eigenvalues can be calculated by following the same steps as in the $q\bar{q}$ case. Indeed, we can consider a tetraquark as a $Q\bar{Q}$ meson, where Q represents the couple of quarks and $\bar{Q}$ the couple of antiquarks (see figure 1), with total “spin” S and relative angular momentum L_{12-34} . The C eigenvectors are those states for which Q and $\bar{Q}$ have opposite charges. So applying the charge conjugation operator to these mesons is the same as exchanging the couple of quarks with the couple of antiquarks. The factors arising from this exchange are the C operator eigenvalues [24]:

$$C = (-1)^{L_{12-34} + S} \quad (13)$$

Tetraquark mesons do not have forbidden J^{PC} combinations because they have more degrees of freedom (in particular they have three different orbital angular momenta) than the normal mesons.

F. Tetraquark states and the Pauli principle

Tetraquarks are composed of two couples of identical fermions, so their states must be antisymmetric for the exchange of the two quarks and the two antiquarks. In this respect it is necessary to study the permutation symmetry

(i.e. the irreducible representations of S_2) of the color, flavor, spin and spatial parts of the wave functions of each subsystem, two quarks and two antiquarks.

Only the singlet color states are physical states, so there are only two color singlets and we write them by underlining their permutation S_2 symmetry, which can be only antisymmetric (A) or symmetric (S):

$$(qq) \text{ in } [\underline{3}]_C \text{ (A) and } (\bar{q}\bar{q}) \text{ in } [3]_C \text{ (A),} \quad (14a)$$

$$(qq) \text{ in } [6]_C \text{ (S) and } (\bar{q}\bar{q}) \text{ in } [\underline{6}]_C \text{ (S).} \quad (14b)$$

Next, we study the permutation symmetry of the spatial part of the two quarks (two antiquarks) states. The permutation symmetry of the spatial part of the couple of quarks and antiquarks is

$$(qq) \text{ with } L_{13} \text{ even (S), } (\bar{q}\bar{q}) \text{ with } L_{24} \text{ even (S)} \quad (15a)$$

$$(qq) \text{ with } L_{13} \text{ odd (A), } (\bar{q}\bar{q}) \text{ with } L_{24} \text{ odd (A)} \quad (15b)$$

$$(qq) \text{ with } L_{13} \text{ even (S), } (\bar{q}\bar{q}) \text{ with } L_{24} \text{ odd (A)} \quad (15c)$$

$$(qq) \text{ with } L_{13} \text{ odd (A), } (\bar{q}\bar{q}) \text{ with } L_{24} \text{ even (S)} \quad (15d)$$

The permutation symmetry of the spatial part derives from the parity of the couple of quarks and antiquarks, which are respectively $P_{qq} = P_q P_q (-1)^{L_{13}} = (-1)^{L_{13}}$ and $P_{\bar{q}\bar{q}} = P_{\bar{q}} P_{\bar{q}} (-1)^{L_{24}} = (-1)^{L_{24}}$.

The permutation symmetry of the $SU(6)_{sf}$ representations for a couple of quarks is written below.

$$[15]_{sf} \text{ (A), which means symmetric spin } (S_{dq} = 1) \text{ and antisymmetric flavor } ([\underline{3}]_f) \\ \text{or antisymmetric spin } (S_{dq} = 0) \text{ and symmetric flavor } ([6]_f) \quad (16a)$$

$$[21]_{sf} \text{ (S), which means symmetric spin } (S_{dq} = 1) \text{ and symmetric flavor } ([6]_f) \\ \text{or antisymmetric spin } (S_{dq} = 0) \text{ and antisymmetric flavor } ([\underline{3}]_f) \quad (16b)$$

The spin-flavor representations for the couple of antiquarks are the conjugate representations $[\overline{15}]_{sf}$ (A) and $[\overline{21}]_{sf}$ (S).

The spatial, flavor, color and spin parts with given permutation symmetry (S_2) must now be arranged together to obtain completely antisymmetric states under the exchange of the two quarks and the two antiquarks. The resulting states are listed in Table I. In this Table we write the color, flavor and spin of the couples of quarks and antiquarks and the corresponding total spin and flavor of the tetraquark states. The total color has been omitted since it is always a singlet.

We want, then, to determine the J^{PC} (where C is obviously intended only for its eigenstates) possible quantum numbers for a tetraquark with a given flavor and spin. The total angular momentum J depends on the values of the three orbital angular momenta L_{13} , L_{24} and L_{12-34} . For obvious reasons, we have chosen to study only the lower value cases, in particular only up to the case that at most one of the three angular momenta is one. In Table II we combine the orbital angular momenta with the spins to obtain the total angular momentum. In Tables III, IV, V and VI we write the possible J^{PC} combinations for every tetraquark with a given flavor and spin.

G. G parity

Charged particles are not eigenstates of C since C takes a positive particle into a negative particle and vice versa. G parity is a generalization of the concept of C parity such that members of an isospin multiplet can each be assigned a good quantum number that would reproduce C for the neutral particle. The G operator is defined as the combination of C and a π rotation around the y axis in the isospin space,

$$G = C\mathcal{R}_y(\pi) = Ce^{i\pi I_2}. \quad (17)$$

The G eigenstates are tetraquark states with flavor charges equal to zero, i.e. strangeness equal to zero in the light mesons case, and their eigenvalues are:

$$G = (-1)^{L_{12-34}+S+I}. \quad (18)$$

The states belonging to $[8] \oplus [10]$ and $[8] \oplus [\overline{10}]$ flavor multiplets are the only exceptions to the validity of Equation (18). Actually a linear combination [25] of these states diagonalizes the G parity.

$$|\psi_G^+ \rangle = \frac{1}{\sqrt{2}}(|[8] \oplus [10] \rangle + |[8] \oplus [\overline{10}] \rangle) \quad (19a)$$

$$|\psi_G^- \rangle = \frac{1}{\sqrt{2}}(|[8] \oplus [10] \rangle - |[8] \oplus [\overline{10}] \rangle) \quad (19b)$$

Table I: Allowed color, flavor and spin tetraquark states.

color $(qq) \otimes (\bar{q}\bar{q})$	L_{13}	L_{24}	S_{dq}	$S_{d\bar{q}}$	flavor $(qq) \otimes (\bar{q}\bar{q})$	S_{tot}	total flavor
$[\bar{3}]_c \otimes [3]_c$	even	even	1	1	$[6]_f \otimes [\bar{6}]_f$	0,1,2	$[1] \oplus [8] \oplus [27]$
$[\bar{3}]_c \otimes [3]_c$	even	even	1	0	$[6]_f \otimes [3]_f$	1	$[8] \oplus [10]$
$[\bar{3}]_c \otimes [3]_c$	even	even	0	1	$[\bar{3}]_f \otimes [\bar{6}]_f$	1	$[8] \oplus [10]$
$[\bar{3}]_c \otimes [3]_c$	even	even	0	0	$[\bar{3}]_f \otimes [3]_f$	0	$[1] \oplus [8]$
$[6]_c \otimes [\bar{6}]_c$	even	even	1	1	$[\bar{3}]_f \otimes [3]_f$	0,1,2	$[1] \oplus [8]$
$[6]_c \otimes [\bar{6}]_c$	even	even	1	0	$[\bar{3}]_f \otimes [\bar{6}]_f$	1	$[8] \oplus [10]$
$[6]_c \otimes [\bar{6}]_c$	even	even	0	1	$[6]_f \otimes [3]_f$	1	$[8] \oplus [10]$
$[6]_c \otimes [\bar{6}]_c$	even	even	0	0	$[6]_f \otimes [\bar{6}]_f$	0	$[1] \oplus [8] \oplus [27]$
$[\bar{3}]_c \otimes [3]_c$	odd	odd	1	1	$[\bar{3}]_f \otimes [3]_f$	0,1,2	$[1] \oplus [8]$
$[\bar{3}]_c \otimes [3]_c$	odd	odd	1	0	$[\bar{3}]_f \otimes [\bar{6}]_f$	1	$[8] \oplus [10]$
$[\bar{3}]_c \otimes [3]_c$	odd	odd	0	1	$[6]_f \otimes [3]_f$	1	$[8] \oplus [10]$
$[\bar{3}]_c \otimes [3]_c$	odd	odd	0	0	$[6]_f \otimes [\bar{6}]_f$	0	$[1] \oplus [8] \oplus [27]$
$[6]_c \otimes [\bar{6}]_c$	odd	odd	1	1	$[6]_f \otimes [\bar{6}]_f$	0,1,2	$[1] \oplus [8] \oplus [27]$
$[6]_c \otimes [\bar{6}]_c$	odd	odd	1	0	$[6]_f \otimes [3]_f$	1	$[8] \oplus [10]$
$[6]_c \otimes [\bar{6}]_c$	odd	odd	0	1	$[\bar{3}]_f \otimes [\bar{6}]_f$	1	$[8] \oplus [10]$
$[6]_c \otimes [\bar{6}]_c$	odd	odd	0	0	$[\bar{3}]_f \otimes [3]_f$	0	$[1] \oplus [8]$
$[\bar{3}]_c \otimes [3]_c$	even	odd	1	1	$[6]_f \otimes [3]_f$	0,1,2	$[8] \oplus [10]$
$[\bar{3}]_c \otimes [3]_c$	even	odd	1	0	$[6]_f \otimes [\bar{6}]_f$	1	$[1] \oplus [8] \oplus [27]$
$[\bar{3}]_c \otimes [3]_c$	even	odd	0	1	$[\bar{3}]_f \otimes [\bar{6}]_f$	1	$[8] \oplus [10]$
$[\bar{3}]_c \otimes [3]_c$	even	odd	0	0	$[\bar{3}]_f \otimes [3]_f$	0	$[1] \oplus [8]$
$[6]_c \otimes [\bar{6}]_c$	even	odd	1	1	$[\bar{3}]_f \otimes [\bar{6}]_f$	0,1,2	$[8] \oplus [10]$
$[6]_c \otimes [\bar{6}]_c$	even	odd	1	0	$[\bar{3}]_f \otimes [3]_f$	1	$[1] \oplus [8]$
$[6]_c \otimes [\bar{6}]_c$	even	odd	0	1	$[6]_f \otimes [\bar{6}]_f$	1	$[1] \oplus [8] \oplus [27]$
$[6]_c \otimes [\bar{6}]_c$	even	odd	0	0	$[6]_f \otimes [3]_f$	0	$[8] \oplus [10]$
$[\bar{3}]_c \otimes [3]_c$	odd	even	1	1	$[\bar{3}]_f \otimes [\bar{6}]_f$	0,1,2	$[8] \oplus [10]$
$[\bar{3}]_c \otimes [3]_c$	odd	even	1	0	$[\bar{3}]_f \otimes [3]_f$	1	$[1] \oplus [8]$
$[\bar{3}]_c \otimes [3]_c$	odd	even	0	1	$[6]_f \otimes [\bar{6}]_f$	1	$[1] \oplus [8] \oplus [27]$
$[\bar{3}]_c \otimes [3]_c$	odd	even	0	0	$[6]_f \otimes [3]_f$	0	$[8] \oplus [10]$
$[6]_c \otimes [\bar{6}]_c$	odd	even	1	1	$[6]_f \otimes [3]_f$	0,1,2	$[8] \oplus [10]$
$[6]_c \otimes [\bar{6}]_c$	odd	even	1	0	$[6]_f \otimes [\bar{6}]_f$	1	$[1] \oplus [8] \oplus [27]$
$[6]_c \otimes [\bar{6}]_c$	odd	even	0	1	$[\bar{3}]_f \otimes [\bar{6}]_f$	1	$[8] \oplus [10]$
$[6]_c \otimes [\bar{6}]_c$	odd	even	0	0	$[\bar{3}]_f \otimes [3]_f$	0	$[1] \oplus [8]$

where $|\psi_G^+\rangle$ and $|\psi_G^-\rangle$ are the G parity eigenvectors with eigenvalues $G = (-1)^{L_{12-34}+S+I+1}$ and $G = (-1)^{L_{12-34}+S+I}$ respectively.

III. THE IACHELLO, MUKHOPADHYAY AND ZHANG MASS FORMULA FOR $q\bar{q}$ MESONS.

In 1991 Iachello, Mukhopadhyay and Zhang developed a mass formula [23, 26] for $q\bar{q}$ mesons, which is a generalization of the Gürsey and Radicati mass formula [27, 28],

$$M^2 = (N_n M_n + N_s M_s)^2 + a \cdot \nu + b \cdot L + c \cdot S + d \cdot J + h \cdot \langle M'^2 \rangle_{ij,i'j'} + i \cdot \langle M''^2 \rangle_{ij,i'j'}, \quad (20)$$

where N_n is the non-strange quark and antiquark number, $M_n = M_u = M_d$ is the non-strange constituent quark mass, N_s is the strange quark and antiquark number, M_s is the strange constituent quark mass, ν is the vibrational quantum number, L is the orbital angular momentum, S the total spin and J the total angular momentum. $\langle M'^2 \rangle_{ij,i'j'}$ and $\langle M''^2 \rangle_{ij,i'j'}$ are two phenomenological terms which act only on the lowest pseudoscalar mesons. Specifically, the first acts on the octet; it encodes the unusually low masses of the bosons of the octet, since they are the eight

Table II: Tetraquark total angular momenta

L_{13}	L_{24}	L_{12-34}	S_{dq}	$S_{d\bar{q}}$	j_{dq}	$j_{d\bar{q}}$	J
0	0	0	0	0	0	0	0
0	0	0	0	1	0	1	1
0	0	0	1	0	1	0	1
0	0	0	1	1	1	1	0,1,2
0	1	0	0	0	0	1	1
0	1	0	0	1	0	0,1,2	0,1,2
0	1	0	1	0	1	1	0,1,2
0	1	0	1	1	1	0,1,2	0,1,2,3
1	0	0	0	0	1	0	1
1	0	0	0	1	1	1	0,1,2
1	0	0	1	0	0,1,2	0	0,1,2
1	0	0	1	1	0,1,2	1	0,1,2,3
0	0	1	0	0	0	0	1
0	0	1	0	1	0	1	0,1,2
0	0	1	1	0	1	0	0,1,2
0	0	1	1	1	1	1	0,1,2,3

Table III: color, flavor, spin (S_{tot}) and J^{PC} for tetraquarks with $L_{13} = L_{24} = L_{12-34} = 0$

color	flavor	S_{dq}	$S_{d\bar{q}}$	j_{dq}	$j_{d\bar{q}}$	S_{tot}	S	J^{PC}
$[\bar{3}] \otimes [3]$	$[1] \oplus [8]$	0	0	0	0	0	0	0^{++}
$[\bar{3}] \otimes [3]$	$[8] \oplus [\bar{10}]$	0	1	0	1	1	1	1^+
$[\bar{3}] \otimes [3]$	$[8] \oplus [10]$	1	0	1	0	1	1	1^+
$[\bar{3}] \otimes [3]$	$[1] \oplus [8] \oplus [27]$	1	1	1	1	0	0	0^{++}
						1	1	1^{+-}
						2	2	2^{++}
$[6] \otimes [\bar{6}]$	$[1] \oplus [8] \oplus [27]$	0	0	0	0	0	0	0^{++}
$[6] \otimes [\bar{6}]$	$[8] \oplus [10]$	0	1	0	1	1	1	1^+
$[6] \otimes [\bar{6}]$	$[8] \oplus [\bar{10}]$	1	0	1	0	1	1	1^+
$[6] \otimes [\bar{6}]$	$[1] \oplus [8]$	1	1	1	1	0	0	0^{++}
						1	1	1^{+-}
						2	2	2^{++}

Goldstone bosons corresponding to the spontaneously broken chiral symmetry group $SU(3)_A$ under which the quark fields transform; the second term acts on the η and η' and relates to the non-negligible $q\bar{q}$ annihilation effects [29] that arise when the lowest mesons are flavor diagonal.

They consider flavor states in the ideal mixing hypothesis, i.e. states with defined number of strange quarks and antiquarks, except for the lowest pseudoscalar nonet. The ideal mixing is essentially a consequence of the OZI rule, introduced by Okubo [30], Zweig [31] and Iizuka [32]. This hypothesis remains to be proved, but it is used by all the authors working on $q\bar{q}$ mesons and also on tetraquarks (see for example Jaffe [6, 7, 20] and Maiani et al. [4]).

Using the updated values for the light $q\bar{q}$ mesons reported by the last PDG [22] (see Tables VII, VIII and IX, see

Table IV: color, flavor, spin (S_{tot}) and J^{PC} for tetraquarks with $L_{13} = L_{12-34} = 0$ and $L_{24} = 1$

color	flavor	S_{dq}	$S_{d\bar{q}}$	j_{dq}	$j_{d\bar{q}}$	S_{tot}	S	J^{PC}
$[3] \otimes [3]$	$[1] \oplus [8]$	0	0	0	1	0	1	1^-
$[3] \otimes [3]$	$[8] \oplus [\overline{10}]$	0	1	0	0	1	0	0^-
					1	1	1	1^-
					2	1	2	2^-
$[\bar{3}] \otimes [3]$	$[1] \oplus [8] \oplus [27]$	1	0	1	1	1	0	0^{-+}
							1	1^{--}
							2	2^{-+}
$[3] \otimes [3]$	$[8] \oplus [10]$	1	1	1	0	0,1,2	1	1^-
					1	0,1,2	0,1,2	$0^-, 1^-, 2^-$
					2	0,1,2	1,2,3	$1^-, 2^-, 3^-$
$[6] \otimes [\bar{6}]$	$[8] \oplus [10]$	0	0	0	1	0	1	1^-
$[6] \otimes [\bar{6}]$	$[1] \oplus [8] \oplus [27]$	0	1	0	0	1	0	0^{-+}
					1	1	1	1^-
					2	1	2	2^-
$[6] \otimes [\bar{6}]$	$[1] \oplus [8]$	1	0	1	1	1	0	0^{-+}
							1	1^{--}
							2	2^{-+}
$[6] \otimes [\bar{6}]$	$[8] \oplus [\overline{10}]$	1	1	1	0	0,1,2	1	1^-
					1	0,1,2	0,1,2	$0^-, 1^-, 2^-$
					2	0,1,2	1,2,3	$1^-, 2^-, 3^-$

Table V: color, flavor, spin (S_{tot}) and J^{PC} for tetraquarks with $L_{24} = L_{12-34} = 0$ and $L_{13} = 1$

color	flavor	S_{dq}	$S_{d\bar{q}}$	j_{dq}	$j_{d\bar{q}}$	S_{tot}	S	J^{PC}
$[3] \otimes [3]$	$[8] \oplus [10]$	0	0	1	0	0	1	1^-
$[3] \otimes [3]$	$[1] \oplus [8] \oplus [27]$	0	1	1	1	1	0	0^{-+}
							1	1^{--}
							2	2^{-+}
$[\bar{3}] \otimes [3]$	$[1] \oplus [8]$	1	0	0	0	1	0	0^{-+}
				1	0	1	1	1^-
				2	0	1	2	2^-
$[3] \otimes [3]$	$[8] \oplus [\overline{10}]$	1	1	0	1	0,1,2	1	1^-
				1	1	0,1,2	0,1,2	$0^-, 1^-, 2^-$
				2	1	0,1,2	1,2,3	$1^-, 2^-, 3^-$
$[6] \otimes [\bar{6}]$	$[1] \oplus [8]$	0	0	1	0	0	1	1^-
$[6] \otimes [\bar{6}]$	$[8] \oplus [\overline{10}]$	0	1	1	1	1	0	0^-
							1	1^-
							2	2^-
$[6] \otimes [\bar{6}]$	$[1] \oplus [8] \oplus [27]$	1	0	0	0	1	0	0^{-+}
				1	0	1	1	1^-
				2	0	1	2	2^-
$[6] \otimes [\bar{6}]$	$[8] \oplus [10]$	1	1	0	1	0,1,2	1	1^-
				1	1	0,1,2	0,1,2	$0^-, 1^-, 2^-$
				2	1	0,1,2	1,2,3	$1^-, 2^-, 3^-$

Table IX: Experimental masses of the K family mesons compared with the theoretical values.

Meson	$M^2(\text{exp.}) \text{ (GeV}^2\text{)}$	$M^2(\text{teo.}) \text{ (GeV}^2\text{)}$	ν	L	S	J^P
(a)						
k(500)	0.24768 ± 0.00001	0.229	0	0	0	0^-
$K^*(892)$	0.8032 ± 0.0004	0.821	0	0	1	1^-
$K_0^*(1430)$	1.99 ± 0.02	1.765	0	1	1	0^+
$K_2^*(1430)$	2.052 ± 0.005	1.939	0	1	1	2^+
$K^*(1680)$	2.95 ± 0.16	2.883	0	2	1	1^-
$K_2(1820)$	3.30 ± 0.09	2.970	0	2	1	2^-
$K_2(1770)$	3.14 ± 0.05	2.891	0	2	0	2^-
$K_3^*(1780)$	3.15 ± 0.04	3.058	0	2	1	3^-
$K^*(1410)$	2.00 ± 0.06	2.210	1	0	1	1^-
(b)						
0^3P_1		1.852	0	1	1	1^+
0^1P_1		1.773	0	1	1	1^+
0^3F_2		4.001	0	3	1	2^+
0^3F_3		4.088	0	3	1	3^+
0^1F_3		4.009	0	3	0	2^+
0^3F_4 ($K_4^*(2045)$)	4.18 ± 0.04	4.176	0	3	1	4^+
1^1S_0 (K(1460))	~ 2.13	2.044	1	0	0	0^-
1^3P_0		3.154	1	1	1	0^+
1^3P_1		3.241	1	1	1	1^+
1^1P_1 ($K_1(1650)$)	2.72 ± 0.17	3.162	1	1	0	1^+
1^3P_2 ($K_2^*(1980)$)	3.89 ± 0.03	3.328	1	1	1	2^+
1^3D_1		4.272	1	2	1	1^-
1^3D_2		4.359	1	2	1	2^-
1^1D_2		4.280	1	2	0	2^-
1^3D_3		4.447	1	2	1	3^-
1^3F_2		5.390	1	3	1	2^+
1^3F_3 ($K_3(2320)$)	5.40 ± 0.11	5.477	1	3	1	3^+

Figure 2: The predicted mesonic masses (solid bars) confronted with the experimental data from the PDG [22], reported with their errors (gray boxes).




Table X: Quantum numbers of the candidate tetraquark nonet

Meson	Mass (GeV)	N_s	$I^G(J^{PC})$	Source
$a_0(980)$	0.9847 ± 0.0012	2	$1^-(0^{++})$	PDG [22]
$f_0(980)$	0.980 ± 0.010	2	$0^+(0^{++})$	PDG [22]
$f_0(600)$	0.478 ± 0.024	0	$0^+(0^{++})$	KLOE [1]
$\kappa(800)$	0.797 ± 0.019	1	$\frac{1}{2}(0^+)$	E791 [38]


As expected, the mesons predicted by the Iachello mass formula reproduce the linear Regge trajectories, representations of $SO(4)$, the linearity of which is satisfied to a high accuracy for light mesons. It is well known that the Regge behaviour [34] can be explained by means of string-like models [35, 36].

IV. THE $qq\bar{q}\bar{q}$ SPECTRUM.

A candidate tetraquark nonet was proposed in the 1970s by Jaffe [6, 7]. This nonet, with quantum numbers $J^{PC} = 0^{++}$, includes the mesons $a_0(980)$, $f_0(980)$, $f_0(600)$ (also called σ meson) and $\kappa(800)$. We hypothesize, as did Jaffe [6, 7, 20], Amsler and Tornqvist [21], Maiani [4] and others, that this nonet is the fundamental tetraquark nonet, with total orbital angular momentum and total spin equal to zero. The candidate tetraquark nonet quantum numbers are presented in Table X, where N_s means the number of strange quarks and antiquarks.

The Iachello, Mukhopadhyay and Zhang mass formula was originally developed for $q\bar{q}$ mesons. In order to describe uncorrelated tetraquark systems by means of an algebraic model one should introduce a new spectrum generating algebra for the spatial part, in this case $U(10)$, since we have nine spatial degrees of freedom. We will not address this difficult problem in this article, but we choose to write the part of the mass formula regarding the internal degrees of freedom in the same way. In $q\bar{q}$ mesons the splitting inside a given flavor multiplet to which is also associated a

Figure 3: Schematic graph of the fundamental tetraquark nonet. The theoretical masses (in MeV), predicted according to Equation (22), are reported below each resonance.



given spin, can be well described by means of the part of the Iachello, Mukhopadhyay and Zhang mass formula that depends only on the numbers of strange and non-strange quarks and antiquarks. It is not necessary, for the purpose of determining the mass splitting of the candidate tetraquark nonet, to calculate the spatial part of the mass formula; we can simply use

$$M^2 = \alpha + (N_n M_n + N_s M_s)^2, \quad (22)$$

where α is a constant that encodes all the spatial and spin dependence of the mass formula, and M_n and M_s are the masses of the constituent quarks (as obtained from an upgrade of the fit of the Iachello, Mukhopadhyay and Zhang mass formula to the new PDG data [22] on $q\bar{q}$ mesons). We determine α by applying Equation (22) to a well-known candidate tetraquark, $a_0(980)$, and in this way we set the energy scale. The value found is

$$\alpha = -1.650 \text{ GeV}^2 \quad (23)$$

With this value of α we predict the masses of the other mesons belonging to the same tetraquark nonet

$$M(\kappa(800)) = 0.726 \text{ GeV} \quad (24)$$

$$M(f_0(600)) = 0.354 \text{ GeV} \quad (25)$$

$$M(f_0(980)) = 0.984 \text{ GeV} \quad (26)$$

The value of $f_0(980)$ agrees very well with the experimental mass reported by the PDG [22], on the contrary our masses of $f_0(600)$ and $\kappa(800)$ are respectively 5 and 4 experimental standard deviations from the values reported in Table X. However, we must remember that the values of the masses $f_0(600)$ and $\kappa(800)$ found by the different experiments are scattered in a range of a few hundreds of MeV around 500 MeV and 800 MeV respectively and the PDG does not report an average mass yet. Thus, new high statistics experiments for the $f_0(600)$ and $\kappa(800)$ are mandatory before reaching any conclusion.

V. DIQUARK-ANTIDIQUARK MODEL

A diquark is a strongly correlated pair of quarks. Since a pair of quarks cannot be a color singlet, the diquark can only be found confined into the hadrons and used as an effective degree of freedom.

Recently many articles have been published regarding the open problem of diquark correlations both in baryons and tetraquarks. Different phenomenological indications for diquarks correlations have been collected over the years as pointed out in Ref. [20] by Jaffe and in Ref. [43] by Selem and Wilczek and references therein, moreover the occurrence of rotational Regge trajectories for baryons with the same slope than the mesonic ones can be explained using a string model [35] of the baryon, where at one end of the string there is a quark in $[3]_c$ color representation and at the other end a diquark in $[\bar{3}]_c$. Recently some papers, relating the physics of the instantons [44, 45, 46, 47], and some calculations in Lattice QCD [48, 49, 50] that support the existence of finite size diquarks as colour antitriplet bound states of two quarks have been published. The diquarks have also been studied in a Coulomb gauge QCD approach [51], that proved their confinement and their well-defined size. One concern is that if diquark correlations are important for exotic states they would already be apparent in the ground state, positive parity nucleon. There is no clear evidence for diquarks in the nucleon, as stated in Refs. [52, 53], and surely completely not for pointlike ones.

From what we have written so far, it is clear that the existence of diquark correlations inside hadrons (and in particular ground state hadrons) is still an open problem. In the meantime, we believe it is not meaningless to study an effective diquark-antidiquark model for tetraquarks. Even if it will be finally found that quarks do not bind together, diquarks as effective degrees of freedom could be useful in hadron spectroscopy in order to correlate many data in terms of a phenomenological model.

A. Classification of the tetraquark states in the diquark-antidiquark model.

We think of the diquark as two correlated quark with no internal spatial excitations, or at least we hypothesize that their internal spatial excitations will be higher in energy than the scale of masses of the resonances we will consider. We describe tetraquark mesons as being composed of a constituent diquark, (qq) , and a constituent antidiquark, $(\bar{q}\bar{q})$. The diquark $SU(3)_c$ color representations are $[\bar{3}]_c$ and $[6]_c$, while the antidiquark ones are $[3]_c$ and $[\bar{6}]_c$, using the standard convention of denoting color and flavor by the dimensions of their representation. As the tetraquark must be a color singlet, the possible diquark-antidiquark color combinations are

$$\text{diquark in } [\bar{3}]_c, \text{ antidiquark in } [3]_c \quad (27a)$$

$$\text{diquark in } [6]_c, \text{ antidiquark in } [\bar{6}]_c \quad (27b)$$

Diquarks (and antidiquarks) are made up of two identical fermions and so they have to satisfy the Pauli principle. Since we consider diquarks with no internal spatial excitations, their color-spin-flavor wave functions must be antisymmetric. This limits the possible representations to being only

$$\text{color in } [\bar{3}] \text{ (AS), spin-flavor in } [21]_{sf} \text{ (S)} \quad (28a)$$

$$\text{color in } [6] \text{ (S), spin-flavor in } [15]_{sf} \text{ (AS)} \quad (28b)$$

This is because we think of the diquark (antidiquark) as two correlated quarks (antiquarks) in an antisymmetric non-excited state. The decomposition of these $SU_{sf}(6)$ representations in terms of $SU(3)_f \otimes SU(2)_s$ is (in the notation [flavor repr., spin])

$$[21]_{sf} = [\bar{3}, 0] \oplus [6, 1] \quad (29a)$$

$$[15]_{sf} = [\bar{3}, 1] \oplus [6, 0] \quad (29b)$$

Using the notation [flavor repr., color repr., spin], the diquark states corresponding to color $[\bar{3}]_c$ and $[6]_c$ respectively, are

$$|[\bar{3}]_f, [\bar{3}]_c, 0\rangle, |[6]_f, [\bar{3}]_c, 1\rangle \quad (30)$$

$$|[\bar{3}]_f, [6]_c, 1\rangle, |[6]_f, [6]_c, 0\rangle \quad (31)$$

The antidiquark states are obtained as the conjugate.

In this paper we will consider only diquarks and antidiquarks in $[\bar{3}]_c$ and $[3]_c$ color representations since like Jaffe [20, 54] or Lichtenberg et al. [55], for example, we expect that color-sextet diquarks will be higher in energy than color-triplet diquarks or even that they will not be bound at all.

Table XII: (continued)

Diquark and antiquark type	total flavor	flavor diquark-antidiquark states with defined n_s ($n_s = 0, 1, 2, 3, 4$)				
		0	1	2	3	4
			$ \frac{3}{2}, I_3, +1\rangle$ $\{n, n\}[\overline{n}, s]$	$ 0, 0, -2\rangle$ $\{s, s\}[\overline{n}, n]$		
$ \overline{6}]_f, [\overline{3}]_c, 1\rangle [\overline{6}]_f, [3]_c, 1\rangle$	$[1] \oplus [8] \oplus [27]$	$ 1, I_3, 0\rangle$; $\{n, n\}\{\overline{n}, n\}$ $ 0, 0, 0\rangle$; $\{n, n\}\{\overline{n}, n\}$ $ 2, I_3, 0\rangle$ $\{n, n\}\{\overline{n}, n\}$	$ \frac{1}{2}, I_3, -1\rangle$; $\{n, s\}\{\overline{n}, n\}$ $ \frac{1}{2}, I_3, +1\rangle$; $\{n, n\}\{\overline{n}, s\}$ $ \frac{3}{2}, I_3, +1\rangle$ $\{n, n\}\{\overline{n}, s\}$ $ \frac{3}{2}, I_3, -1\rangle$ $\{n, s\}\{\overline{n}, n\}$	$ 0, 0, 0\rangle$; $\{n, s\}\{\overline{n}, s\}$ $ 1, I_3, 0\rangle$; $\{n, s\}\{\overline{n}, s\}$ $ 1, I_3, +2\rangle$ $\{n, n\}\{\overline{s}, s\}$ $ 1, I_3, -2\rangle$ $\{s, s\}\{\overline{n}, n\}$	$ \frac{1}{2}, I_3, -1\rangle$; $\{s, s\}\{\overline{n}, s\}$ $ \frac{1}{2}, I_3, +1\rangle$ $\{n, s\}\{\overline{s}, s\}$	$ 0, 0, 0\rangle$ $\{s, s\}\{\overline{s}, s\}$
(end of table)						

Table XIII: Spectroscopic classification and diquark content of tetraquarks states with $L_{12-34} = 0$ in the diquark-antidiquark limit.

$^{2S+1}L_J(J^{PC})$	Diquark and antidiquark type
$^1S_0(0^{++})$	$ \overline{3}]_f, [\overline{3}]_c, 0\rangle [\overline{3}]_f, [3]_c, 0\rangle$; $ \overline{6}]_f, [\overline{3}]_c, 1\rangle [\overline{6}]_f, [3]_c, 1\rangle$
$^3S_1(1^+)$	$ \overline{3}]_f, [\overline{3}]_c, 0\rangle [\overline{6}]_f, [3]_c, 1\rangle$; $ \overline{6}]_f, [\overline{3}]_c, 1\rangle [\overline{3}]_f, [3]_c, 0\rangle$
$^3S_1(1^{+-})$	$ \overline{6}]_f, [\overline{3}]_c, 1\rangle [\overline{6}]_f, [3]_c, 1\rangle$
$^5S_2(2^{++})$	$ \overline{6}]_f, [\overline{3}]_c, 1\rangle [\overline{6}]_f, [3]_c, 1\rangle$

Table XIV: As in Table XIII but for $L_{12-34} = 1$

$^{2S+1}L_J(J^{PC})$	Diquark and antidiquark type
$^1P_1(1^{--})$	$ \overline{3}]_f, [\overline{3}]_c, 0\rangle [\overline{3}]_f, [3]_c, 0\rangle$; $ \overline{6}]_f, [\overline{3}]_c, 1\rangle [\overline{6}]_f, [3]_c, 1\rangle$
$^3P_2(2^-)$	$ \overline{3}]_f, [\overline{3}]_c, 0\rangle [\overline{6}]_f, [3]_c, 1\rangle$; $ \overline{6}]_f, [\overline{3}]_c, 1\rangle [\overline{3}]_f, [3]_c, 0\rangle$
$^3P_2(2^{+-})$	$ \overline{6}]_f, [\overline{3}]_c, 1\rangle [\overline{6}]_f, [3]_c, 1\rangle$
$^3P_1(1^-)$	$ \overline{3}]_f, [\overline{3}]_c, 0\rangle [\overline{6}]_f, [3]_c, 1\rangle$; $ \overline{6}]_f, [\overline{3}]_c, 1\rangle [\overline{3}]_f, [3]_c, 0\rangle$
$^3P_1(1^{+-})$	$ \overline{6}]_f, [\overline{3}]_c, 1\rangle [\overline{6}]_f, [3]_c, 1\rangle$
$^3P_0(0^-)$	$ \overline{3}]_f, [\overline{3}]_c, 0\rangle [\overline{6}]_f, [3]_c, 1\rangle$; $ \overline{6}]_f, [\overline{3}]_c, 1\rangle [\overline{3}]_f, [3]_c, 0\rangle$
$^3P_0(0^{+-})$	$ \overline{6}]_f, [\overline{3}]_c, 1\rangle [\overline{6}]_f, [3]_c, 1\rangle$
$^5P_3(3^{--})$	$ \overline{6}]_f, [\overline{3}]_c, 1\rangle [\overline{6}]_f, [3]_c, 1\rangle$
$^5P_2(2^{--})$	$ \overline{6}]_f, [\overline{3}]_c, 1\rangle [\overline{6}]_f, [3]_c, 1\rangle$
$^5P_1(1^{--})$	$ \overline{6}]_f, [\overline{3}]_c, 1\rangle [\overline{6}]_f, [3]_c, 1\rangle$

How to read Tables XII, XIII, XIV, XV can be explained by examples. In Table XII we can read the diquark content of the states belonging to a given flavor multiplet. For example, as we can read from the first line of Table XII, the nine states belonging to the flavor nonet are made up of two scalar diquarks. In particular the state $|0, 0, 0\rangle$ contains the $[n, n]$ diquark and the $[\overline{n}, \overline{n}]$ antidiquark. Tables XIII, XIV, XV show for a given J^{PC} which diquark-antidiquark type content is possible and also which J^{PC} quantum numbers can be assigned to a given diquark-antidiquark state. For example, as indicated in the first line of Table XIII, the only possible tetraquarks with $^1S_0(0^{++})$ quantum numbers are those containing two scalar diquarks or two vector diquarks (which correspond respectively to the flavor nonet and the flavor 36-plet, as we can see in Table XII).

Table XV: As in Table XIII but for $L_{12-34} = 2$

$^{2S+1}L_J(J^{PC})$	Diquark and antidiquark type
$^1D_2(2^{++})$	$ \bar{3}_f, \bar{3}_c, 0\rangle \bar{3}_f, \bar{3}_c, 0\rangle;$ $ \bar{6}_f, \bar{3}_c, 1\rangle \bar{6}_f, \bar{3}_c, 1\rangle$
$^3D_3(3^+)$	$ \bar{3}_f, \bar{3}_c, 0\rangle \bar{6}_f, \bar{3}_c, 1\rangle;$ $ \bar{6}_f, \bar{3}_c, 1\rangle \bar{3}_f, \bar{3}_c, 0\rangle$
$^3D_3(3^{+-})$	$ \bar{6}_f, \bar{3}_c, 1\rangle \bar{6}_f, \bar{3}_c, 1\rangle$
$^3D_2(2^+)$	$ \bar{3}_f, \bar{3}_c, 0\rangle \bar{6}_f, \bar{3}_c, 1\rangle;$ $ \bar{6}_f, \bar{3}_c, 1\rangle \bar{3}_f, \bar{3}_c, 0\rangle$
$^3D_2(2^{+-})$	$ \bar{6}_f, \bar{3}_c, 1\rangle \bar{6}_f, \bar{3}_c, 1\rangle$
$^3D_1(1^+)$	$ \bar{3}_f, \bar{3}_c, 0\rangle \bar{6}_f, \bar{3}_c, 1\rangle;$ $ \bar{6}_f, \bar{3}_c, 1\rangle \bar{3}_f, \bar{3}_c, 0\rangle$
$^3D_1(1^{+-})$	$ \bar{6}_f, \bar{3}_c, 1\rangle \bar{6}_f, \bar{3}_c, 1\rangle$
$^5D_4(4^{++})$	$ \bar{6}_f, \bar{3}_c, 1\rangle \bar{6}_f, \bar{3}_c, 1\rangle$
$^5D_3(3^{++})$	$ \bar{6}_f, \bar{3}_c, 1\rangle \bar{6}_f, \bar{3}_c, 1\rangle$
$^5D_2(2^{++})$	$ \bar{6}_f, \bar{3}_c, 1\rangle \bar{6}_f, \bar{3}_c, 1\rangle$
$^5D_1(1^{++})$	$ \bar{6}_f, \bar{3}_c, 1\rangle \bar{6}_f, \bar{3}_c, 1\rangle$
$^5D_0(0^{++})$	$ \bar{6}_f, \bar{3}_c, 1\rangle \bar{6}_f, \bar{3}_c, 1\rangle$

Table XVI: Quantum numbers of the candidate tetraquark nonet. $\kappa(800)$ corresponds to $[n, n] \overline{[n, s]}$ and also to its conjugate.

Meson	Mass (GeV)	Diquark content	$I^G(J^{PC})$	Source
$a_0(980)$	0.9847 ± 0.0012	$[n, s] \overline{[n, s]}$	$1^-(0^{++})$	PDG [22]
$f_0(980)$	0.980 ± 0.010	$[n, s] \overline{[n, s]}$	$0^+(0^{++})$	PDG [22]
$f_0(600)$	0.478 ± 0.024	$[n, n] \overline{[n, n]}$	$0^+(0^{++})$	KLOE [1]
$\kappa(800)$	0.797 ± 0.019	$[n, n] \overline{[n, s]}$	$\frac{1}{2}(0^+)$	E791 [38]

B. The tetraquark nonet spectrum in the diquark-antidiquark model.

We describe diquark-antidiquark tetraquark configurations by using $U(4) \otimes SU(3)_f \otimes SU(2)_s \otimes SU(3)_c$ as spectrum generating algebra, by analogy with what was done by Iachello et al. [23, 26] for the normal mesons. In a string model [35, 36] the slopes of these trajectories depend only on the color representation of the constituent particles. Thus the slope of Regge trajectories of tetraquarks made up of a diquark in $\bar{3}_c$ and an antidiquark in 3_c is the same as the slope of Regge trajectories of $q\bar{q}$ mesons.

For the tetraquark in the diquark-antidiquark model, we can use the mass formula developed by Iachello et al. [23, 26] for the normal mesons, but replacing the masses of the quark and the antiquark with those of the diquark and the antidiquark:

$$M^2 = (M_{qq} + M_{\bar{q}\bar{q}})^2 + a \cdot n + b \cdot L_{12-34} + c \cdot S_{tot} + d \cdot J, \quad (32)$$

where M_{qq} and $M_{\bar{q}\bar{q}}$ are the diquark and antidiquark masses, n is a vibrational quantum number, L_{12-34} the relative orbital angular momentum, S_{tot} the total spin and J the total angular momentum.

We need, then, to determine the diquark masses. This can be done by fitting the mass formula (32) with the mass values of the tetraquark candidate nonet [63] $a_0(980)$, $f_0(980)$, $f_0(600)$ and $\kappa(800)$. Following Jaffe's arguments [6, 20, 54], the candidate tetraquark nonet is the fundamental tetraquark multiplet and it contains the lighter diquarks, i.e. scalar diquarks.

The masses of the scalar diquarks resulting from the fit are:

$$M_{[n, n]} = 0.275 \text{ GeV}, \quad M_{[n, s]} = 0.492 \text{ GeV} \quad (33)$$

The masses of the candidate tetraquark nonet obtained from the fit[64] are:

$$M_{a_0(980)} = M_{f_0(980)} = 0.984 \text{ GeV} \quad (34a)$$

$$M_{f_0(600)} = 0.550 \text{ GeV} \quad (34b)$$

$$M_{\kappa(800)} = 0.767 \text{ GeV}. \quad (34c)$$

The masses of $a_0(980)$ and $f_0(980)$ agree with the experimental values reported by the PDG [22], $\kappa(800)$ and $f_0(600)$ are respectively within 2 and 3 experimental standard deviations from the values reported in Table XVI.

The value of $f_0(600)$ is similar to the value recently found by Mathur, Nilmani and others [56] in a tetraquark model with Lattice QCD.

VI. SUMMARY, CONCLUSIONS AND OUTLOOK

In this work, we have constructed a complete classification scheme of the tetraquark states in terms of $SU(6)_{sf}$ spin-flavor multiplets, and their flavor and spin content in terms of $SU(3)_f$ and $SU(2)_s$ states. Moreover, we have discussed the permutation symmetry properties of both the spin-flavor and orbital parts of the qq and $\bar{q}\bar{q}$ subsystems. In order to obtain the total wave function, the spin-flavor part has been combined with the color and orbital contributions in such a way that the total tetraquark wave function is a color singlet and is antisymmetric under the interchange of the two quarks and the two antiquarks. This classification scheme is general and complete, and may be helpful for experimental, CQM and lattice QCD studies. In particular, the constructed basis for tetraquark states will enable the eigenvalue problem to be solved for a definite dynamical model.

As an application, we have calculated the mass spectrum of the candidate tetraquark nonet, adapting to the tetraquark case the Iachello, Mukhopadhyay and Zhang [23, 26] mass formula, developed originally for the qq mesons. We have considered only the part of this formula that gives the splitting inside a given multiplet.

The predicted tetraquark states in the low energy range are much more numerous than the experimental candidate tetraquarks. So, if the tetraquark model is correct, we must solve the problem of the missing tetraquark resonances.

The introduction of the diquark-antidiquark model, in section V, helps us to drastically reduce the number of predicted tetraquark states. In fact the allowed states in this model are only a small subset of the states in the uncorrelated model. Nevertheless this cut is not sufficient and the remaining tetraquark resonances are still too numerous. Thus, we need another explanation for the missing resonances.

If it is heavy enough, a $qq\bar{q}\bar{q}$ meson will be unstable against decay into two $q\bar{q}$ mesons. The $qq\bar{q}\bar{q}$ state simply falls apart, or dissociates [6, 7]. Thus, we can deduce that if a given $qq\bar{q}\bar{q}$ state is above threshold for decay into a “dissociation” channel, it is very broad into that channel and the higher the energy of the resonance is the broader the phase space is.

The great width of most $qq\bar{q}\bar{q}$ mesons will account for their experimental elusiveness, thus making it difficult to establish their resonant character at all. The $f_0(600)$ provides a clear example of this. Many higher-mass $qq\bar{q}\bar{q}$ states not only may be as broad and confusing as the $f_0(600)$, but also will probably occur in channels with substantial inelastic background obscuring their resonant behaviour.

As an alternative to the tetraquark hypothesis, the possibility was considered that $a_0(980)$ and $f_0(980)$ may be $K - \bar{K}$ bound states, kept together by hadron exchange forces, the same that bind nucleons in the nuclei, color singlet remnants of the confining color forces (hence the name $K - \bar{K}$ molecules [57] used in this connection). If they are indeed $K - \bar{K}$ molecules, scalar mesons do not need to make a complete $SU(3)_f$ multiplet so that this idea would be consistent with the lack of evidence of $f_0(600)$ and $\kappa(800)$. However, if the existence of these particle were confirmed, it would be very hard to consider either of them as a $\pi - \pi$ or $\pi - K$ molecule, since the latter particles would in any case lie considerably higher than the respective thresholds. We see that the existence or absence of these light scalars is crucial in assessing the nature of $a_0(980)$ and $f_0(980)$. From this point of view, the recent observations of $f_0(600)$ and $\kappa(800)$ in D non-leptonic decays at Fermilab[38] and in the $\pi\pi$ spectrum in $\phi \rightarrow \pi^0\pi^0\gamma$ at Frascati [38, 58] have considerably reinforced the hypothesis of a full tetraquark nonet with inverted spectrum. The experimental situation and the latest results concerning $f_0(600)$ and $\kappa(800)$ are summarized in Refs. [3, 59, 60] and references therein. New high-statistics experiments dedicated to these resonances are important in order to confirm or refute their existence.

Appendix A: THE $qq\bar{q}\bar{q}$ FLAVOR STATES

In this appendix we write the $qq\bar{q}\bar{q}$ flavor states explicitly in terms of the states of the single quarks and antiquarks (the color states may be easily obtained from the flavor ones by using the replacements $u \leftrightarrow r$, $d \leftrightarrow g$ and $s \leftrightarrow b$). These states are calculated by using the $SU(3)$ Clebsch-Gordan coefficients [61, 62] in the De Swart [62] phase convention.

A generic flavor state is expressed by $[[R]I, I_3, Y >$, where $[R]$ indicates the $SU(3)_f$ representation, I the isospin quantum number, I_3 its third component and Y the hypercharge. The single quark states are written in short as u , d and s , where

$$u = |[3]\frac{1}{2}, -\frac{1}{2}, +\frac{1}{3} >$$

$$d = |[3]\frac{1}{2}, +\frac{1}{2}, +\frac{1}{3} >$$

$$s = |[3]0, 0, -\frac{2}{3} > .$$

In this appendix the tetraquark states are written in the $(qq)(\bar{q}\bar{q})$ configuration; thus, in addition to the representation $[R]$ to which the state belongs, we also show the representations $[R']$ and $[R'']$ respectively of the two quarks and the two antiquarks. In short, a tetraquark state will be written as $[[R]I, I_3, Y >_{R' R''}$. The states in the $q\bar{q}q\bar{q}$ configuration are a linear combination of those in the $(qq)(\bar{q}\bar{q})$ configuration.

1. The tetraquark flavor states in the $(qq)(\bar{q}\bar{q})$ configuration

$$|[1]0, 0, 0 >_{\bar{3}3} = \frac{1}{2\sqrt{3}}(su\bar{s}\bar{u} - su\bar{u}\bar{s} - us\bar{s}\bar{u} + us\bar{u}\bar{s} - ds\bar{d}\bar{s} + ds\bar{s}\bar{d} + sdd\bar{s} - sd\bar{d}\bar{s} + ud\bar{u}\bar{d} - udd\bar{u} - du\bar{u}\bar{d} + dud\bar{u}) \quad (A1)$$

$$|[1]0, 0, 0 >_{6\bar{6}} = \frac{1}{2\sqrt{6}}(2uu\bar{u}\bar{u} - ud\bar{u}\bar{d} - udd\bar{u} - du\bar{u}\bar{d} - dud\bar{u} + 2ddd\bar{d} - us\bar{u}\bar{s} - us\bar{s}\bar{u} - su\bar{u}\bar{s} - su\bar{s}\bar{u} + ds\bar{d}\bar{s} + ds\bar{s}\bar{d} + sdd\bar{s} + sd\bar{s}\bar{d} + 2ss\bar{s}\bar{s}) \quad (A2)$$

$$|[8]1, +1, 0 >_{\bar{3}3} = \frac{1}{2}(-sud\bar{s} + su\bar{s}\bar{d} + us\bar{d}\bar{s} - us\bar{s}\bar{d}) \quad (A3)$$

$$|[8]1, 0, 0 >_{\bar{3}3} = \frac{1}{2\sqrt{2}}(-sd\bar{d}\bar{s} - su\bar{u}\bar{s} + sd\bar{s}\bar{d} + su\bar{s}\bar{u} + ds\bar{d}\bar{s} + us\bar{u}\bar{s} - ds\bar{s}\bar{d} - us\bar{s}\bar{u}) \quad (A4)$$

$$|[8]1, -1, 0 >_{\bar{3}3} = \frac{1}{2}(-ds\bar{s}\bar{u} + ds\bar{u}\bar{s} + sd\bar{s}\bar{u} - sd\bar{u}\bar{s}) \quad (A5)$$

$$|[8]\frac{1}{2}, +\frac{1}{2}, +1 >_{\bar{3}3} = \frac{1}{2}(-udd\bar{s} + ud\bar{s}\bar{d} + dud\bar{s} - du\bar{s}\bar{d}) \quad (A6)$$

$$|[8]\frac{1}{2}, -\frac{1}{2}, +1 >_{\bar{3}3} = \frac{1}{2}(-ud\bar{u}\bar{s} + ud\bar{s}\bar{u} + du\bar{u}\bar{s} - du\bar{s}\bar{u}) \quad (A7)$$

$$|[8]\frac{1}{2}, +\frac{1}{2}, -1 >_{\bar{3}3} = \frac{1}{2}(su\bar{u}\bar{d} - su\bar{d}\bar{u} - us\bar{u}\bar{d} + us\bar{d}\bar{u}) \quad (A8)$$

$$|[8]\frac{1}{2}, -\frac{1}{2}, -1 >_{\bar{3}3} = \frac{1}{2}(-ds\bar{u}\bar{d} + ds\bar{d}\bar{u} + sd\bar{u}\bar{d} - sdd\bar{u}) \quad (A9)$$

$$|[8]0, 0, 0 >_{\bar{3}3} = \frac{1}{2\sqrt{6}}(-su\bar{s}\bar{u} + su\bar{u}\bar{s} + us\bar{s}\bar{u} - us\bar{u}\bar{s} + ds\bar{d}\bar{s} - ds\bar{s}\bar{d} - sdd\bar{s} + sd\bar{s}\bar{d} + 2ud\bar{u}\bar{d} - 2udd\bar{u} - 2du\bar{u}\bar{d} + 2dud\bar{u}) \quad (A10)$$

$$|[8]1, +1, 0\rangle_{63} = \frac{1}{2\sqrt{3}}(-2uu\bar{u}\bar{d} + 2u\bar{u}\bar{d}\bar{u} - us\bar{d}\bar{s} + us\bar{s}\bar{d} - su\bar{d}\bar{s} + su\bar{s}\bar{d}) \quad (\text{A11})$$

$$|[8]1, 0, 0\rangle_{63} = \frac{1}{2\sqrt{6}}(-2du\bar{u}\bar{d} - 2u\bar{u}\bar{d}\bar{d} + 2du\bar{d}\bar{u} + 2u\bar{d}\bar{d}\bar{u} + ds\bar{s}\bar{d} + us\bar{s}\bar{u} - ds\bar{d}\bar{s} - us\bar{u}\bar{s} - s\bar{d}\bar{d}\bar{s} - su\bar{u}\bar{s} + s\bar{d}\bar{s}\bar{d} + su\bar{s}\bar{u}) \quad (\text{A12})$$

$$|[8]1, -1, 0\rangle_{63} = \frac{1}{2\sqrt{3}}(2dd\bar{d}\bar{u} - 2d\bar{d}\bar{u}\bar{d} - ds\bar{u}\bar{s} + ds\bar{s}\bar{u} - s\bar{d}\bar{u}\bar{s} + s\bar{d}\bar{s}\bar{u}) \quad (\text{A13})$$

$$|[8]\frac{1}{2}, +\frac{1}{2}, +1\rangle_{63} = \frac{1}{2\sqrt{3}}(2uu\bar{s}\bar{u} - 2uu\bar{u}\bar{s} + u\bar{d}\bar{s}\bar{s} - u\bar{s}\bar{d}\bar{s} + du\bar{d}\bar{s} - du\bar{s}\bar{d}) \quad (\text{A14})$$

$$|[8]\frac{1}{2}, -\frac{1}{2}, +1\rangle_{63} = \frac{1}{2\sqrt{3}}(2dd\bar{d}\bar{s} - 2d\bar{d}\bar{s}\bar{d} + du\bar{s}\bar{u} + u\bar{d}\bar{s}\bar{u} - du\bar{u}\bar{s} - u\bar{d}\bar{u}\bar{s}) \quad (\text{A15})$$

$$|[8]\frac{1}{2}, +\frac{1}{2}, -1\rangle_{63} = \frac{1}{2\sqrt{3}}(2ss\bar{s}\bar{d} - 2s\bar{s}\bar{d}\bar{s} + su\bar{d}\bar{u} - su\bar{u}\bar{d} + us\bar{d}\bar{u} - us\bar{u}\bar{d}) \quad (\text{A16})$$

$$|[8]\frac{1}{2}, -\frac{1}{2}, -1\rangle_{63} = \frac{1}{2\sqrt{3}}(2ss\bar{s}\bar{u} - 2s\bar{s}\bar{u}\bar{s} + s\bar{d}\bar{d}\bar{u} - s\bar{d}\bar{u}\bar{d} + ds\bar{d}\bar{u} - ds\bar{u}\bar{d}) \quad (\text{A17})$$

$$|[8]0, 0, 0\rangle_{63} = \frac{1}{2\sqrt{2}}(su\bar{s}\bar{u} - su\bar{u}\bar{s} + us\bar{s}\bar{u} - us\bar{u}\bar{s} + ds\bar{d}\bar{s} - ds\bar{s}\bar{d} + s\bar{d}\bar{d}\bar{s} - s\bar{d}\bar{s}\bar{d}) \quad (\text{A18})$$

$$|[8]1, +1, 0\rangle_{\bar{3}6} = \frac{1}{2\sqrt{3}}(2u\bar{d}\bar{d}\bar{d} - 2du\bar{d}\bar{d} + us\bar{d}\bar{s} + us\bar{s}\bar{d} - su\bar{d}\bar{s} - su\bar{s}\bar{d}) \quad (\text{A19})$$

$$|[8]1, 0, 0\rangle_{\bar{3}6} = \frac{1}{2\sqrt{6}}(-2du\bar{u}\bar{d} + 2u\bar{u}\bar{d}\bar{d} - 2du\bar{d}\bar{u} + 2u\bar{d}\bar{d}\bar{u} + ds\bar{s}\bar{d} + us\bar{s}\bar{u} + ds\bar{d}\bar{s} + us\bar{u}\bar{s} - s\bar{d}\bar{d}\bar{s} - su\bar{u}\bar{s} - s\bar{d}\bar{s}\bar{d} - su\bar{s}\bar{u}) \quad (\text{A20})$$

$$|[8]1, -1, 0\rangle_{\bar{3}6} = \frac{1}{2\sqrt{3}}(2u\bar{d}\bar{u}\bar{u} - 2du\bar{u}\bar{u} + ds\bar{u}\bar{s} + ds\bar{s}\bar{u} - s\bar{d}\bar{u}\bar{s} - s\bar{d}\bar{s}\bar{u}) \quad (\text{A21})$$

$$|[8]\frac{1}{2}, +\frac{1}{2}, +1\rangle_{\bar{3}6} = \frac{1}{2\sqrt{3}}(2us\bar{s}\bar{s} - 2su\bar{s}\bar{s} + u\bar{d}\bar{s}\bar{s} + u\bar{s}\bar{d}\bar{s} - du\bar{d}\bar{s} - du\bar{s}\bar{d}) \quad (\text{A22})$$

$$|[8]\frac{1}{2}, -\frac{1}{2}, +1\rangle_{\bar{3}6} = \frac{1}{2\sqrt{3}}(2ds\bar{s}\bar{s} - 2s\bar{d}\bar{s}\bar{s} + u\bar{d}\bar{u}\bar{s} - du\bar{u}\bar{s} + u\bar{d}\bar{s}\bar{u} - du\bar{s}\bar{u}) \quad (\text{A23})$$

$$|[8]\frac{1}{2}, +\frac{1}{2}, -1\rangle_{\bar{3}6} = \frac{1}{2\sqrt{3}}(2s\bar{d}\bar{d}\bar{d} - 2ds\bar{d}\bar{d} - su\bar{u}\bar{d} + us\bar{u}\bar{d} - su\bar{d}\bar{u} - us\bar{d}\bar{u}) \quad (\text{A24})$$

$$|[8]_{\frac{1}{2}, -\frac{1}{2}, -1} >_{\bar{3}\bar{6}} = \frac{1}{2\sqrt{3}}(2us\bar{u}\bar{u} - 2su\bar{u}\bar{u} + sd\bar{u}\bar{d} + sdd\bar{u} + ds\bar{u}\bar{d} - ds\bar{d}\bar{u}) \quad (\text{A25})$$

$$|[8]0, 0, 0 >_{\bar{3}\bar{6}} = \frac{1}{2\sqrt{2}}(-su\bar{s}\bar{u} - su\bar{u}\bar{s} + us\bar{s}\bar{u} + us\bar{u}\bar{s} - ds\bar{d}\bar{s} + ds\bar{s}\bar{d} + sdd\bar{s} + sd\bar{s}\bar{d}) \quad (\text{A26})$$

$$|[8]1, +1, 0 >_{\bar{6}\bar{6}} = \frac{1}{2\sqrt{5}}(2uu\bar{u}\bar{d} + 2uu\bar{d}\bar{u} - 2udd\bar{d} - 2du\bar{d}\bar{d} + us\bar{d}\bar{s} - us\bar{s}\bar{d} - sud\bar{s} - su\bar{s}\bar{d}) \quad (\text{A27})$$

$$|[8]1, 0, 0 >_{\bar{6}\bar{6}} = \frac{1}{2\sqrt{10}}(4uu\bar{u}\bar{u} - 4ddd\bar{d} - ds\bar{d}\bar{s} - us\bar{u}\bar{s} + ds\bar{s}\bar{d} - us\bar{s}\bar{u} - sdd\bar{s} - su\bar{u}\bar{s} - sd\bar{s}\bar{d} - su\bar{s}\bar{u}) \quad (\text{A28})$$

$$|[8]1, -1, 0 >_{\bar{6}\bar{6}} = \frac{1}{2\sqrt{5}}(2du\bar{u}\bar{u} + 2ud\bar{u}\bar{u} - 2dd\bar{u}\bar{d} - 2dd\bar{d}\bar{u} + ds\bar{u}\bar{s} - ds\bar{s}\bar{u} - sd\bar{u}\bar{s} - sd\bar{s}\bar{u}) \quad (\text{A29})$$

$$|[8]_{\frac{1}{2}, +\frac{1}{2}, +1} >_{\bar{6}\bar{6}} = \frac{1}{2\sqrt{5}}(2uu\bar{u}\bar{s} + 2uu\bar{s}\bar{u} - 2us\bar{s}\bar{s} - 2su\bar{s}\bar{s} + udd\bar{s} - ud\bar{s}\bar{d} - dud\bar{s} - du\bar{s}\bar{d}) \quad (\text{A30})$$

$$|[8]_{\frac{1}{2}, -\frac{1}{2}, +1} >_{\bar{6}\bar{6}} = \frac{1}{2\sqrt{5}}(-2ds\bar{s}\bar{s} - 2sd\bar{s}\bar{s} - 2dd\bar{s}\bar{d} - 2dd\bar{d}\bar{s} + ud\bar{u}\bar{s} + du\bar{u}\bar{s} + ud\bar{s}\bar{u} + du\bar{s}\bar{u}) \quad (\text{A31})$$

$$|[8]_{\frac{1}{2}, +\frac{1}{2}, -1} >_{\bar{6}\bar{6}} = \frac{1}{2\sqrt{5}}(-2sdd\bar{d} - 2dsdd\bar{d} - 2ssd\bar{s} - 2ss\bar{s}\bar{d} + su\bar{u}\bar{d} + us\bar{u}\bar{d} + sud\bar{u} + us\bar{d}\bar{u}) \quad (\text{A32})$$

$$|[8]_{\frac{1}{2}, -\frac{1}{2}, -1} >_{\bar{6}\bar{6}} = \frac{1}{2\sqrt{5}}(2us\bar{u}\bar{u} + 2su\bar{u}\bar{u} - 2ss\bar{u}\bar{s} - 2ss\bar{s}\bar{u} + sd\bar{u}\bar{d} - sdd\bar{u} - ds\bar{u}\bar{d} - ds\bar{d}\bar{u}) \quad (\text{A33})$$

$$|[8]0, 0, 0 >_{\bar{6}\bar{6}} = \frac{1}{2\sqrt{30}}(4uu\bar{u}\bar{u} + 4ddd\bar{d} - 8ss\bar{s}\bar{s} + su\bar{s}\bar{u} + su\bar{u}\bar{s} + us\bar{s}\bar{u} + us\bar{u}\bar{s} - ds\bar{d}\bar{s} - ds\bar{s}\bar{d} - sdd\bar{s} - sd\bar{s}\bar{d} - 2ud\bar{u}\bar{d} - 2udd\bar{u} - 2du\bar{u}\bar{d} - 2du\bar{d}\bar{u}) \quad (\text{A34})$$

$$|[10]_{\frac{3}{2}, +\frac{3}{2}, +1} >_{\bar{6}3} = \frac{1}{\sqrt{2}}(uud\bar{s} - uu\bar{s}\bar{d}) \quad (\text{A35})$$

$$|[10]_{\frac{3}{2}, +\frac{1}{2}, +1} >_{\bar{6}3} = \frac{1}{\sqrt{6}}(uu\bar{u}\bar{s} - uu\bar{s}\bar{u} + dud\bar{s} - du\bar{s}\bar{d} + udd\bar{s} - ud\bar{s}\bar{d}) \quad (\text{A36})$$

$$|[10]_{\frac{3}{2}, -\frac{1}{2}, +1} >_{\bar{6}3} = \frac{1}{\sqrt{6}}(dd\bar{d}\bar{s} - dd\bar{s}\bar{d} + du\bar{u}\bar{s} - du\bar{s}\bar{u} + ud\bar{u}\bar{s} - ud\bar{s}\bar{u}) \quad (\text{A37})$$

$$|[10]_{\frac{3}{2}, -\frac{3}{2}, +1} >_{\bar{6}3} = \frac{1}{\sqrt{2}}(dd\bar{u}\bar{s} - dd\bar{s}\bar{u}) \quad (\text{A38})$$

$$|[10]1, +1, 0 >_{\bar{6}3} = \frac{1}{\sqrt{6}}(uud\bar{u} - uu\bar{u}\bar{d} + us\bar{d}\bar{s} - us\bar{s}\bar{d} + sud\bar{s} - su\bar{s}\bar{d}) \quad (\text{A39})$$

$$|[10]1, 0, 0 \rangle_{63} = \frac{1}{2\sqrt{3}}(-du\bar{u}\bar{d} - ud\bar{u}\bar{d} + du\bar{d}\bar{u} + udd\bar{u} + ds\bar{s}\bar{d} - us\bar{s}\bar{u} + ds\bar{d}\bar{s} + us\bar{u}\bar{s} + sdd\bar{s} + su\bar{u}\bar{s} - sd\bar{s}\bar{d} - su\bar{s}\bar{u}) \quad (\text{A40})$$

$$|[10]1, -1, 0 \rangle_{63} = \frac{1}{\sqrt{6}}(ddd\bar{u} - dd\bar{u}\bar{d} + ds\bar{u}\bar{s} - ds\bar{s}\bar{u} + sd\bar{u}\bar{s} - sd\bar{s}\bar{u}) \quad (\text{A41})$$

$$|[10]\frac{1}{2}, +\frac{1}{2}, -1 \rangle_{63} = \frac{1}{\sqrt{6}}(usd\bar{u} - us\bar{u}\bar{d} + ss\bar{d}\bar{s} - ss\bar{s}\bar{d} + sud\bar{u} - su\bar{u}\bar{d}) \quad (\text{A42})$$

$$|[10]\frac{1}{2}, -\frac{1}{2}, -1 \rangle_{63} = \frac{1}{\sqrt{6}}(dsd\bar{u} - ds\bar{u}\bar{d} + ss\bar{u}\bar{s} - ss\bar{s}\bar{u} + sdd\bar{u} - sd\bar{u}\bar{d}) \quad (\text{A43})$$

$$|[10]0, 0, -2 \rangle_{63} = \frac{1}{\sqrt{2}}(ssd\bar{u} - ss\bar{u}\bar{d}) \quad (\text{A44})$$

$$|[\overline{10}]\frac{3}{2}, +\frac{3}{2}, -1 \rangle_{\overline{3}\overline{6}} = \frac{1}{\sqrt{2}}(usd\bar{d} - sud\bar{d}) \quad (\text{A45})$$

$$|[\overline{10}]\frac{3}{2}, +\frac{1}{2}, -1 \rangle_{\overline{3}\overline{6}} = \frac{1}{\sqrt{6}}(dsd\bar{d} - sdd\bar{d} + us\bar{u}\bar{d} - su\bar{u}\bar{d} + usd\bar{u} - sud\bar{u}) \quad (\text{A46})$$

$$|[\overline{10}]\frac{3}{2}, -\frac{1}{2}, -1 \rangle_{\overline{3}\overline{6}} = \frac{1}{\sqrt{6}}(ds\bar{u}\bar{d} - sd\bar{u}\bar{d} + ds\bar{d}\bar{u} - sdd\bar{u} + us\bar{u}\bar{u} - su\bar{u}\bar{u}) \quad (\text{A47})$$

$$|[\overline{10}]\frac{3}{2}, -\frac{3}{2}, -1 \rangle_{\overline{3}\overline{6}} = \frac{1}{\sqrt{2}}(ds\bar{u}\bar{u} - sd\bar{u}\bar{u}) \quad (\text{A48})$$

$$|[\overline{10}]1, +1, 0 \rangle_{\overline{3}\overline{6}} = \frac{1}{\sqrt{6}}(du\bar{d}\bar{d} - udd\bar{d} + us\bar{d}\bar{s} + us\bar{s}\bar{d} - sud\bar{s} - su\bar{s}\bar{d}) \quad (\text{A49})$$

$$|[\overline{10}]1, 0, 0 \rangle_{\overline{3}\overline{6}} = \frac{1}{2\sqrt{3}}(du\bar{u}\bar{d} - ud\bar{u}\bar{d} + du\bar{d}\bar{u} - udd\bar{u} + ds\bar{s}\bar{d} + us\bar{s}\bar{u} + ds\bar{d}\bar{s} + us\bar{u}\bar{s} - sdd\bar{s} - su\bar{u}\bar{s} - sd\bar{s}\bar{d} - su\bar{s}\bar{u}) \quad (\text{A50})$$

$$|[\overline{10}]1, -1, 0 \rangle_{\overline{3}\overline{6}} = \frac{1}{\sqrt{6}}(du\bar{u}\bar{u} - ud\bar{u}\bar{u} + ds\bar{u}\bar{s} + ds\bar{s}\bar{u} - sd\bar{u}\bar{s} - sd\bar{s}\bar{u}) \quad (\text{A51})$$

$$|[\overline{10}]\frac{1}{2}, +\frac{1}{2}, +1 \rangle_{\overline{3}\overline{6}} = \frac{1}{\sqrt{6}}(us\bar{s}\bar{s} - su\bar{s}\bar{s} + du\bar{d}\bar{s} - udd\bar{s} + du\bar{s}\bar{d} - ud\bar{s}\bar{d}) \quad (\text{A52})$$

$$|[\overline{10}]\frac{1}{2}, -\frac{1}{2}, +1 \rangle_{\overline{3}\overline{6}} = \frac{1}{\sqrt{6}}(ds\bar{s}\bar{s} - sd\bar{s}\bar{s} + du\bar{u}\bar{s} - ud\bar{u}\bar{s} + du\bar{s}\bar{u} - ud\bar{s}\bar{u}) \quad (\text{A53})$$

$$|[\overline{10}]0, 0, +2 \rangle_{\overline{3}\overline{6}} = \frac{1}{\sqrt{2}}(du\bar{s}\bar{s} - ud\bar{s}\bar{s}) \quad (\text{A54})$$

$$|[27]2, +2, 0 \rangle_{6\overline{6}} = (uud\bar{d}) \quad (\text{A55})$$

$$|[27]2, +1, 0 >_{6\bar{6}} = \frac{1}{2}(du\bar{d}\bar{d} + udd\bar{d} + uu\bar{u}\bar{d} + uu\bar{d}\bar{u}) \quad (\text{A56})$$

$$|[27]2, 0, 0 >_{6\bar{6}} = \frac{1}{\sqrt{6}}(uu\bar{u}\bar{u} + ud\bar{u}\bar{d} + dd\bar{d}\bar{d} + du\bar{d}\bar{u} + du\bar{u}\bar{d} + udd\bar{u}) \quad (\text{A57})$$

$$|[27]2, -1, 0 >_{6\bar{6}} = \frac{1}{2}(du\bar{u}\bar{u} + ud\bar{u}\bar{u} + dd\bar{u}\bar{d} + dd\bar{d}\bar{u}) \quad (\text{A58})$$

$$|[27]2, -2, 0 >_{6\bar{6}} = (dd\bar{u}\bar{u}) \quad (\text{A59})$$

$$|[27]\frac{3}{2}, +\frac{3}{2}, +1 >_{6\bar{6}} = \frac{1}{\sqrt{2}}(uud\bar{s} + uu\bar{s}\bar{d}) \quad (\text{A60})$$

$$|[27]\frac{3}{2}, +\frac{1}{2}, +1 >_{6\bar{6}} = \frac{1}{\sqrt{6}}(uu\bar{u}\bar{s} + uu\bar{s}\bar{u} + du\bar{d}\bar{s} + du\bar{s}\bar{d} + udd\bar{s} + ud\bar{s}\bar{d}) \quad (\text{A61})$$

$$|[27]\frac{3}{2}, -\frac{1}{2}, +1 >_{6\bar{6}} = \frac{1}{\sqrt{6}}(dd\bar{d}\bar{s} + dd\bar{s}\bar{d} + du\bar{u}\bar{s} + du\bar{s}\bar{u} + ud\bar{u}\bar{s} + ud\bar{s}\bar{u}) \quad (\text{A62})$$

$$|[27]\frac{3}{2}, -\frac{3}{2}, +1 >_{6\bar{6}} = \frac{1}{\sqrt{2}}(dd\bar{u}\bar{s} + dd\bar{s}\bar{u}) \quad (\text{A63})$$

$$|[27]\frac{3}{2}, +\frac{3}{2}, -1 >_{6\bar{6}} = \frac{1}{\sqrt{2}}(us\bar{d}\bar{d} + su\bar{d}\bar{d}) \quad (\text{A64})$$

$$|[27]\frac{3}{2}, +\frac{1}{2}, -1 >_{6\bar{6}} = \frac{1}{\sqrt{6}}(ds\bar{d}\bar{d} + sdd\bar{d} + us\bar{u}\bar{d} + su\bar{u}\bar{d} + us\bar{d}\bar{u} + su\bar{d}\bar{u}) \quad (\text{A65})$$

$$|[27]\frac{3}{2}, -\frac{1}{2}, -1 >_{6\bar{6}} = \frac{1}{\sqrt{6}}(ds\bar{u}\bar{d} + sd\bar{u}\bar{d} + ds\bar{d}\bar{u} + sdd\bar{u} + us\bar{u}\bar{u} + su\bar{u}\bar{u}) \quad (\text{A66})$$

$$|[27]\frac{3}{2}, -\frac{3}{2}, -1 >_{6\bar{6}} = \frac{1}{\sqrt{2}}(ds\bar{u}\bar{u} + sd\bar{u}\bar{u}) \quad (\text{A67})$$

$$|[27]1, +1, +2 >_{6\bar{6}} = (uu\bar{s}\bar{s}) \quad (\text{A68})$$

$$|[27]1, 0, +2 >_{6\bar{6}} = \frac{1}{\sqrt{2}}(du\bar{s}\bar{s} + ud\bar{s}\bar{s}) \quad (\text{A69})$$

$$|[27]1, -1, +2 >_{6\bar{6}} = (dd\bar{s}\bar{s}) \quad (\text{A70})$$

$$|[27]1, +1, -2 >_{6\bar{6}} = (ss\bar{d}\bar{d}) \quad (\text{A71})$$

$$|[27]1, 0, -2 >_{6\bar{6}} = \frac{1}{\sqrt{2}}(ss\bar{u}\bar{d} + ss\bar{d}\bar{u}) \quad (\text{A72})$$

$$|[27]1, -1, -2 >_{6\bar{6}} = (ss\bar{u}\bar{u}) \quad (\text{A73})$$

$$|[27]1, +1, 0 >_{6\bar{6}} = \frac{1}{2\sqrt{5}}(uu\bar{u}\bar{d} + uu\bar{d}\bar{u} - udd\bar{d} - du\bar{d}\bar{d} + 2us\bar{d}\bar{s} + 2us\bar{s}\bar{d} + 2su\bar{d}\bar{s} + 2su\bar{s}\bar{d}) \quad (\text{A74})$$

$$|[27]1, 0, 0 >_{\bar{6}\bar{6}} = \frac{1}{\sqrt{10}}(uu\bar{u}\bar{u} - ddd\bar{d} + ds\bar{d}\bar{s} + us\bar{u}\bar{s} + ds\bar{s}\bar{d} + us\bar{s}\bar{u} + sdd\bar{s} + su\bar{u}\bar{s} + sd\bar{s}\bar{d} + su\bar{s}\bar{u}) \quad (\text{A75})$$

$$|[27]1, -1, 0 >_{\bar{6}\bar{6}} = \frac{1}{2\sqrt{5}}(du\bar{u}\bar{u} + ud\bar{u}\bar{u} - dd\bar{u}\bar{d} - dd\bar{d}\bar{u} + 2ds\bar{u}\bar{s} + 2ds\bar{s}\bar{u} + 2sd\bar{u}\bar{s} + 2sd\bar{s}\bar{u}) \quad (\text{A76})$$

$$|[27]\frac{1}{2}, +\frac{1}{2}, +1 >_{\bar{6}\bar{6}} = \frac{1}{\sqrt{30}}(2uu\bar{u}\bar{s} + 2uu\bar{s}\bar{u} + 3us\bar{s}\bar{s} + 3su\bar{s}\bar{s} - udd\bar{s} - ud\bar{s}\bar{d} - du\bar{d}\bar{s} - du\bar{s}\bar{d}) \quad (\text{A77})$$

$$|[27]\frac{1}{2}, -\frac{1}{2}, +1 >_{\bar{6}\bar{6}} = \frac{1}{\sqrt{30}}(-2dd\bar{s}\bar{d} - 2dd\bar{d}\bar{s} + 3ds\bar{s}\bar{s} + 3sd\bar{s}\bar{s} + ud\bar{u}\bar{s} + du\bar{u}\bar{s} + ud\bar{s}\bar{u} + du\bar{s}\bar{u}) \quad (\text{A78})$$

$$|[27]\frac{1}{2}, +\frac{1}{2}, -1 >_{\bar{6}\bar{6}} = \frac{1}{\sqrt{30}}(-2sdd\bar{d} - 2ds\bar{d}\bar{d} + 3ss\bar{d}\bar{s} + 3ss\bar{s}\bar{d} + su\bar{u}\bar{d} + us\bar{u}\bar{d} + su\bar{d}\bar{u} + us\bar{d}\bar{u}) \quad (\text{A79})$$

$$|[27]\frac{1}{2}, -\frac{1}{2}, -1 >_{\bar{6}\bar{6}} = \frac{1}{\sqrt{30}}(2us\bar{u}\bar{u} + 2su\bar{u}\bar{u} + 3ss\bar{u}\bar{s} + 3ss\bar{s}\bar{u} - sd\bar{u}\bar{d} - sdd\bar{u} - ds\bar{u}\bar{d} - ds\bar{d}\bar{u}) \quad (\text{A80})$$

$$|[27]0, 0, 0 >_{\bar{6}\bar{6}} = \frac{1}{2\sqrt{30}}(2uu\bar{u}\bar{u} + 2ddd\bar{d} + 6ss\bar{s}\bar{s} + 3su\bar{s}\bar{u} + 3su\bar{u}\bar{s} + 3us\bar{s}\bar{u} + 3us\bar{u}\bar{s} - 3ds\bar{d}\bar{s} - 3ds\bar{s}\bar{d} - 3sdd\bar{s} - 3sd\bar{s}\bar{d} - ud\bar{u}\bar{d} - udd\bar{u} - du\bar{u}\bar{d} - du\bar{d}\bar{u}) \quad (\text{A81})$$

2. The $qq\bar{q}\bar{q}$ states in the “ideal mixing” hypothesis

In the “ideal mixing” hypothesis, the flavor states of the tetraquarks are a superposition of the states written in section (A 1) in such a way to have defined strange quark and antiquark numbers. The notation used for the ideally mixed states is $|\text{strange quark number}, I, I_3, Y >_{R' R''}$. Clearly the only states that can be mixed are those with the same quantum numbers (i.e. same isospin and same hypercharge). Only the mixed states are written below.

$$|n_S = 2, 0, 0, 0 >_{\bar{3}3} = \frac{1}{\sqrt{3}}(\sqrt{2}[8]0, 0, 0 >_{\bar{3}3} + |[1]0, 0, 0 >_{\bar{3}3}) = \frac{1}{2\sqrt{2}}(-su\bar{s}\bar{u} + su\bar{u}\bar{s} + us\bar{s}\bar{u} - us\bar{u}\bar{s} + ds\bar{d}\bar{s} - ds\bar{s}\bar{d} - sdd\bar{s} + sd\bar{s}\bar{d}) \quad (\text{A82})$$

$$|n_S = 0, 0, 0, 0 >_{\bar{3}3} = \frac{1}{\sqrt{3}}([8]0, 0, 0 >_{\bar{3}3} - \sqrt{2}[1]0, 0, 0 >_{\bar{3}3}) = \frac{1}{2}(ud\bar{u}\bar{d} - udd\bar{u} - du\bar{u}\bar{d} + du\bar{d}\bar{u}) \quad (\text{A83})$$

$$|n_S = 2, 1, +1, 0 >_{\bar{3}\bar{6}} = \frac{1}{\sqrt{3}}(\sqrt{2}[\overline{10}]1, +1, 0 >_{\bar{3}\bar{6}} + [8]1, +1, 0 >_{\bar{3}\bar{6}}) = \frac{1}{2}(us\bar{d}\bar{s} + us\bar{s}\bar{d} - su\bar{d}\bar{s} - su\bar{s}\bar{d}) \quad (\text{A84})$$

$$|n_S = 2, 1, 0, 0 >_{\bar{3}\bar{6}} = \frac{1}{2\sqrt{2}}(ds\bar{d}\bar{s} + us\bar{u}\bar{s} + ds\bar{s}\bar{d} + us\bar{s}\bar{u} - sdd\bar{s} - su\bar{u}\bar{s} - sd\bar{s}\bar{d} - su\bar{s}\bar{u}) \quad (\text{A85})$$

$$|n_S = 2, 1, -1, 0 >_{\bar{3}\bar{6}} = \frac{1}{2}(ds\bar{u}\bar{s} + ds\bar{s}\bar{u} - sd\bar{u}\bar{s} - sd\bar{s}\bar{u}) \quad (\text{A86})$$

$$\begin{aligned}
|n_S = 0, 1, +1, 0 >_{\bar{3}\bar{6}} &= \frac{1}{\sqrt{3}}(|[\bar{10}]1, +1, 0 >_{\bar{3}\bar{6}} - \sqrt{2}[8]0, 0, 0 >_{\bar{3}\bar{6}}) = \\
&= \frac{1}{\sqrt{2}}(dud\bar{d} - ud\bar{d})
\end{aligned} \tag{A87}$$

$$|n_S = 2, 1, 0, 0 >_{\bar{3}\bar{6}} = \frac{1}{2}(du\bar{u}\bar{d} + du\bar{d}\bar{u} - ud\bar{u}\bar{d} - ud\bar{d}\bar{u}) \tag{A88}$$

$$|n_S = 0, 1, -1, 0 >_{\bar{3}\bar{6}} = \frac{1}{\sqrt{2}}(du\bar{u}\bar{u} - ud\bar{u}\bar{u}) \tag{A89}$$

$$\begin{aligned}
|n_S = 3, \frac{1}{2}, +\frac{1}{2}, +1 >_{\bar{3}\bar{6}} &= \frac{1}{\sqrt{3}}(|[\bar{10}]\frac{1}{2}, +\frac{1}{2}, +1 >_{\bar{3}\bar{6}} + \sqrt{2}[8]\frac{1}{2}, +\frac{1}{2}, +1 >_{\bar{3}\bar{6}}) = \\
&= \frac{1}{\sqrt{2}}(us\bar{s}\bar{s} - su\bar{s}\bar{s})
\end{aligned} \tag{A90}$$

$$|n_S = 3, \frac{1}{2}, -\frac{1}{2}, +1 >_{\bar{3}\bar{6}} = \frac{1}{\sqrt{2}}(ds\bar{s}\bar{s} - sd\bar{s}\bar{s}) \tag{A91}$$

$$\begin{aligned}
|n_S = 1, \frac{1}{2}, +\frac{1}{2}, +1 >_{\bar{3}\bar{6}} &= \frac{1}{\sqrt{3}}(-\sqrt{2}[\bar{10}]\frac{1}{2}, +\frac{1}{2}, +1 >_{\bar{3}\bar{6}} + [8]\frac{1}{2}, +\frac{1}{2}, +1 >_{\bar{3}\bar{6}}) = \\
&= \frac{1}{2}(udd\bar{s} - dud\bar{s} + ud\bar{s}\bar{d} - du\bar{s}\bar{d})
\end{aligned} \tag{A92}$$

$$|n_S = 1, \frac{1}{2}, -\frac{1}{2}, +1 >_{\bar{3}\bar{6}} = \frac{1}{2}(ud\bar{u}\bar{s} - du\bar{u}\bar{s} + ud\bar{s}\bar{u} - du\bar{s}\bar{u}) \tag{A93}$$

$$\begin{aligned}
|n_S = 2, 1, +1, 0 >_{63} &= \frac{1}{\sqrt{3}}(-\sqrt{2}[10]1, +1, 0 >_{63} + [8]1, +1, 0 >_{63}) = \\
&= \frac{1}{2}(us\bar{s}\bar{d} - us\bar{d}\bar{s} - sud\bar{s} + su\bar{s}\bar{d})
\end{aligned} \tag{A94}$$

$$|n_S = 2, 1, 0, 0 >_{63} = \frac{1}{2\sqrt{2}}(-ds\bar{d}\bar{s} - us\bar{u}\bar{s} + ds\bar{s}\bar{d} + us\bar{s}\bar{u} - sdd\bar{s} - su\bar{u}\bar{s} + sd\bar{s}\bar{d} + su\bar{s}\bar{u}) \tag{A95}$$

$$|n_S = 2, 1, -1, 0 >_{63} = \frac{1}{2}(-ds\bar{u}\bar{s} + ds\bar{s}\bar{u} - sd\bar{u}\bar{s} + sd\bar{s}\bar{u}) \tag{A96}$$

$$\begin{aligned}
|n_S = 0, 1, +1, 0 >_{63} &= \frac{1}{\sqrt{3}}([10]1, +1, 0 >_{63} + \sqrt{2}[8]1, +1, 0 >_{63}) = \\
&= \frac{1}{\sqrt{2}}(uud\bar{u} - uu\bar{u}\bar{d})
\end{aligned} \tag{A97}$$

$$|n_S = 0, 1, 0, 0 >_{63} = \frac{1}{2}(-du\bar{u}\bar{d} + du\bar{d}\bar{u} - ud\bar{u}\bar{d} + ud\bar{d}\bar{u}) \tag{A98}$$

$$|n_S = 0, 1, -1, 0 >_{63} = \frac{1}{\sqrt{2}}(dd\bar{d}\bar{u} - dd\bar{u}\bar{d}) \tag{A99}$$

$$\begin{aligned}
|n_S = 3, \frac{1}{2}, +\frac{1}{2}, -1 >_{63} &= \frac{1}{\sqrt{3}}([10]\frac{1}{2}, +\frac{1}{2}, -1 >_{63} - \sqrt{2}[8]\frac{1}{2}, +\frac{1}{2}, -1 >_{63}) = \\
&= \frac{1}{\sqrt{2}}(ss\bar{d}\bar{s} - ss\bar{s}\bar{d})
\end{aligned} \tag{A100}$$

$$|n_S = 3, \frac{1}{2}, -\frac{1}{2}, -1 >_{63} = \frac{1}{\sqrt{2}}(ss\bar{u}\bar{s} - ss\bar{s}\bar{u}) \tag{A101}$$

$$|n_S = 1, \frac{1}{2}, +\frac{1}{2}, -1 \rangle_{63} = \frac{1}{\sqrt{3}}(\sqrt{2}[10]_{\frac{1}{2}, +\frac{1}{2}, -1} \rangle_{63} + |[8]_{\frac{1}{2}, +\frac{1}{2}, -1} \rangle_{63}) = \frac{1}{2}(sud\bar{u} - su\bar{u}\bar{d} + us\bar{d}\bar{u} - us\bar{u}\bar{d}) \quad (\text{A102})$$

$$|n_S = 1, \frac{1}{2}, -\frac{1}{2}, -1 \rangle_{63} = \frac{1}{2}(sdd\bar{u} - sd\bar{u}\bar{d} + ds\bar{d}\bar{u} - ds\bar{u}\bar{d}) \quad (\text{A103})$$

$$|n_S = 2, 1, +1, 0 \rangle_{6\bar{6}} = \frac{1}{\sqrt{5}}(2|[27]1, +1, 0 \rangle_{6\bar{6}} - |[8]1, +1, 0 \rangle_{6\bar{6}}) = \frac{1}{2}(us\bar{s}\bar{d} + us\bar{d}\bar{s} + su\bar{d}\bar{s} + su\bar{s}\bar{d}) \quad (\text{A104})$$

$$|n_S = 2, 1, 0, 0 \rangle_{6\bar{6}} = \frac{1}{2\sqrt{2}}(ds\bar{d}\bar{s} + us\bar{u}\bar{s} + ds\bar{s}\bar{d} + us\bar{s}\bar{u} + sdd\bar{s} + su\bar{u}\bar{s} + sd\bar{s}\bar{d} + su\bar{s}\bar{u}) \quad (\text{A105})$$

$$|n_S = 2, 1, -1, 0 \rangle_{6\bar{6}} = \frac{1}{2}(ds\bar{u}\bar{s} + ds\bar{s}\bar{u} + sd\bar{u}\bar{s} + sd\bar{s}\bar{u}) \quad (\text{A106})$$

$$|n_S = 0, 1, +1, 0 \rangle_{6\bar{6}} = \frac{1}{\sqrt{5}}(-|[27]1, +1, 0 \rangle_{6\bar{6}} - 2|[8]1, +1, 0 \rangle_{6\bar{6}}) = \frac{1}{2}(udd\bar{d} + dud\bar{d} - uu\bar{u}\bar{d} - uu\bar{d}\bar{u}) \quad (\text{A107})$$

$$|n_S = 0, 1, 0, 0 \rangle_{6\bar{6}} = \frac{1}{\sqrt{2}}(ddd\bar{d} - uu\bar{u}\bar{u}) \quad (\text{A108})$$

$$|n_S = 0, 1, -1, 0 \rangle_{6\bar{6}} = \frac{1}{2}(dd\bar{u}\bar{d} + dd\bar{d}\bar{u} - du\bar{u}\bar{u} - ud\bar{u}\bar{u}) \quad (\text{A109})$$

$$|n_S = 3, \frac{1}{2}, +\frac{1}{2}, -1 \rangle_{6\bar{6}} = \frac{1}{\sqrt{5}}(\sqrt{3}[27]_{\frac{1}{2}, +\frac{1}{2}, -1} \rangle_{6\bar{6}} - \sqrt{2}[8]_{\frac{1}{2}, +\frac{1}{2}, -1} \rangle_{6\bar{6}}) = \frac{1}{\sqrt{2}}(ss\bar{d}\bar{s} + ss\bar{s}\bar{d}) \quad (\text{A110})$$

$$|n_S = 3, \frac{1}{2}, -\frac{1}{2}, -1 \rangle_{6\bar{6}} = \frac{1}{\sqrt{2}}(ss\bar{u}\bar{s} + ss\bar{s}\bar{u}) \quad (\text{A111})$$

$$|n_S = 1, \frac{1}{2}, +\frac{1}{2}, -1 \rangle_{6\bar{6}} = \frac{1}{\sqrt{5}}(\sqrt{2}[27]_{\frac{1}{2}, +\frac{1}{2}, -1} \rangle_{6\bar{6}} + \sqrt{3}[8]_{\frac{1}{2}, +\frac{1}{2}, -1} \rangle_{6\bar{6}}) = \frac{1}{2\sqrt{3}}(sud\bar{u} + su\bar{u}\bar{d} + us\bar{d}\bar{u} + us\bar{u}\bar{d} - 2sdd\bar{d} - 2ds\bar{d}\bar{d}) \quad (\text{A112})$$

$$|n_S = 1, \frac{1}{2}, -\frac{1}{2}, -1 \rangle_{6\bar{6}} = \frac{1}{2\sqrt{3}}(2us\bar{u}\bar{u} + 2su\bar{u}\bar{u} - sdd\bar{u} - sd\bar{u}\bar{d} - ds\bar{d}\bar{u} - ds\bar{u}\bar{d}) \quad (\text{A113})$$

$$|n_S = 3, \frac{1}{2}, +\frac{1}{2}, +1 \rangle_{6\bar{6}} = \frac{1}{\sqrt{5}}(\sqrt{3}[27]_{\frac{1}{2}, +\frac{1}{2}, +1} \rangle_{6\bar{6}} - \sqrt{2}[8]_{\frac{1}{2}, +\frac{1}{2}, +1} \rangle_{6\bar{6}}) = \frac{1}{\sqrt{2}}(us\bar{s}\bar{s} + su\bar{s}\bar{s}) \quad (\text{A114})$$

$$|n_S = 3, \frac{1}{2}, -\frac{1}{2}, +1 \rangle_{6\bar{6}} = \frac{1}{\sqrt{2}}(ds\bar{s}\bar{s} + sd\bar{s}\bar{s}) \quad (\text{A115})$$

$$|n_S = 1, \frac{1}{2}, +\frac{1}{2}, +1 \rangle_{6\bar{6}} = \frac{1}{\sqrt{5}}(\sqrt{2}[27]_{\frac{1}{2}, +\frac{1}{2}, +1} \rangle_{6\bar{6}} + \sqrt{3}[8]_{\frac{1}{2}, +\frac{1}{2}, +1} \rangle_{6\bar{6}}) = \frac{1}{2\sqrt{3}}(2uu\bar{u}\bar{s} + 2uu\bar{s}\bar{u} - udd\bar{s} - dud\bar{s} - ud\bar{s}\bar{d} - du\bar{s}\bar{d}) \quad (\text{A116})$$

$$|n_S = 1, \frac{1}{2}, -\frac{1}{2}, +1 \rangle_{6\bar{6}} = \frac{1}{2\sqrt{3}}(ud\bar{u}\bar{s} + du\bar{u}\bar{s} + ud\bar{s}\bar{u} + du\bar{s}\bar{u} - 2dd\bar{d}\bar{s} - 2dd\bar{s}\bar{d}) \quad (\text{A117})$$

$$|n_S = 4, 0, 0, 0 \rangle_{6\bar{6}} = \frac{\sqrt{3}}{\sqrt{10}}|[27]0, 0, 0 \rangle_{6\bar{6}} - \frac{2\sqrt{2}}{\sqrt{15}}|[8]0, 0, 0 \rangle_{6\bar{6}} + \frac{1}{\sqrt{6}}|[1]0, 0, 0 \rangle_{6\bar{6}} = ss\bar{s}\bar{s} \quad (\text{A118})$$

$$|n_S = 2, 0, 0, 0 \rangle_{6\bar{6}} = -\frac{\sqrt{3}}{\sqrt{5}}|[27]0, 0, 0 \rangle_{6\bar{6}} - \frac{1}{\sqrt{15}}|[8]0, 0, 0 \rangle_{6\bar{6}} + \frac{1}{\sqrt{3}}|[1]0, 0, 0 \rangle_{6\bar{6}} = \frac{1}{2\sqrt{2}}(-us\bar{u}\bar{s} - us\bar{s}\bar{u} - su\bar{u}\bar{s} - su\bar{s}\bar{u} + ds\bar{d}\bar{s} + ds\bar{s}\bar{d} + sdd\bar{s} + sd\bar{s}\bar{d}) \quad (\text{A119})$$

$$|n_S = 0, 0, 0, 0 \rangle_{6\bar{6}} = -\frac{1}{\sqrt{10}}|[27]0, 0, 0 \rangle_{6\bar{6}} - \frac{\sqrt{2}}{\sqrt{5}}|[8]0, 0, 0 \rangle_{6\bar{6}} - \frac{1}{\sqrt{2}}|[1]0, 0, 0 \rangle_{6\bar{6}} = \frac{1}{2\sqrt{3}}(ud\bar{u}\bar{d} + du\bar{u}\bar{d} + udd\bar{u} + dud\bar{u} - 2uu\bar{u}\bar{u} - 2dd\bar{d}\bar{d}) \quad (\text{A120})$$

Appendix B: THE TETRAQUARK SPIN STATES

In this appendix we write the spin states of the tetraquarks in terms of the spins of the single quarks and antiquarks. These states are calculated by using the SU(2) Clebsch-Gordan coefficients.

A generic spin state is expressed by $|[R] S, S_z \rangle$, where $[R]$ indicates the SU(2)_S representation, S the spin quantum number and S_z its third component. The single quark states are written in short as: $\uparrow \equiv |[2] \frac{1}{2} + \frac{1}{2} \rangle$ and $\downarrow \equiv |[2] \frac{1}{2} - \frac{1}{2} \rangle$.

In this appendix the tetraquark states are written in the $(qq)(\bar{q}\bar{q})$ configuration; thus, in addition to the representation $[R]$ to which the state belongs, we also show the representations $[R']$ and $[R'']$ respectively of the two quarks and the two antiquarks. In short a tetraquark state will be written as $|[R] S, S_z \rangle_{R' R''}$.

1. Tetraquark spin states in the $(qq)(\bar{q}\bar{q})$ configuration

$$|[1] 0 0 \rangle_{11} = \frac{1}{2}(\uparrow\downarrow\uparrow\downarrow + \downarrow\uparrow\downarrow\uparrow - \uparrow\downarrow\downarrow\uparrow - \downarrow\uparrow\uparrow\downarrow) \quad (\text{B1})$$

$$|[1] 0 0 \rangle_{33} = \frac{1}{2\sqrt{3}}(2 \uparrow\uparrow\downarrow\downarrow + 2 \downarrow\downarrow\uparrow\uparrow + \uparrow\downarrow\uparrow\downarrow + \downarrow\uparrow\downarrow\uparrow + \uparrow\downarrow\downarrow\uparrow + \downarrow\uparrow\uparrow\downarrow) \quad (\text{B2})$$

$$|[3] 1 +1 \rangle_{31} = \frac{1}{\sqrt{2}}(\uparrow\uparrow\uparrow\downarrow - \uparrow\uparrow\downarrow\uparrow) \quad (\text{B3})$$

$$|[3] 1 0 \rangle_{31} = \frac{1}{2}(\uparrow\downarrow\uparrow\downarrow + \downarrow\uparrow\uparrow\downarrow - \uparrow\downarrow\downarrow\uparrow - \downarrow\uparrow\downarrow\uparrow) \quad (\text{B4})$$

$$|[3] 1 -1 \rangle_{31} = \frac{1}{\sqrt{2}}(\downarrow\downarrow\uparrow\downarrow - \downarrow\downarrow\downarrow\uparrow) \quad (\text{B5})$$

$$|[3] 1 +1 \rangle_{13} = \frac{1}{\sqrt{2}}(\uparrow\downarrow\uparrow\uparrow - \downarrow\uparrow\uparrow\uparrow) \quad (\text{B6})$$

$$|[3] 1 0 \rangle_{13} = \frac{1}{2}(\uparrow\downarrow\uparrow\downarrow + \uparrow\downarrow\downarrow\uparrow - \downarrow\uparrow\uparrow\downarrow - \downarrow\uparrow\downarrow\uparrow) \quad (\text{B7})$$

$$|[3] 1 - 1 >_{13} = \frac{1}{\sqrt{2}}(\uparrow\downarrow\downarrow\downarrow - \downarrow\uparrow\downarrow\downarrow) \quad (\text{B8})$$

$$|[3] 1 + 1 >_{33} = \frac{1}{2}(\uparrow\uparrow\uparrow\downarrow + \uparrow\uparrow\downarrow\uparrow - \uparrow\downarrow\uparrow\uparrow - \downarrow\uparrow\uparrow\uparrow) \quad (\text{B9})$$

$$|[3] 1 0 >_{33} = \frac{1}{\sqrt{2}}(\uparrow\uparrow\downarrow\downarrow - \downarrow\downarrow\uparrow\uparrow) \quad (\text{B10})$$

$$|[3] 1 - 1 >_{33} = \frac{1}{2}(-\downarrow\downarrow\uparrow\downarrow - \downarrow\downarrow\downarrow\uparrow + \uparrow\downarrow\downarrow\downarrow + \downarrow\uparrow\downarrow\downarrow) \quad (\text{B11})$$

$$|[5] 2 + 2 >_{33} = \uparrow\uparrow\uparrow\uparrow \quad (\text{B12})$$

$$|[5] 2 + 1 >_{33} = \frac{1}{2}(\uparrow\uparrow\uparrow\downarrow + \uparrow\uparrow\downarrow\uparrow + \uparrow\downarrow\uparrow\uparrow + \downarrow\uparrow\uparrow\uparrow) \quad (\text{B13})$$

$$|[5] 2 0 >_{33} = \frac{1}{\sqrt{6}}(\uparrow\uparrow\downarrow\downarrow + \downarrow\downarrow\uparrow\uparrow + \uparrow\downarrow\uparrow\downarrow + \uparrow\downarrow\downarrow\uparrow + \downarrow\uparrow\uparrow\downarrow + \downarrow\downarrow\uparrow\downarrow) \quad (\text{B14})$$

$$|[5] 2 - 1 >_{33} = \frac{1}{2}(\downarrow\downarrow\uparrow\downarrow + \downarrow\downarrow\downarrow\uparrow + \uparrow\downarrow\downarrow\downarrow + \downarrow\uparrow\downarrow\downarrow) \quad (\text{B15})$$

$$|[5] 2 - 2 >_{33} = \downarrow\downarrow\downarrow\downarrow \quad (\text{B16})$$

-
- [1] A. Aloisio et al. (KLOE), Phys. Lett. **B537**, 21 (2002), hep-ex/0204013.
 - [2] E. M. Aitala et al. (E791), Phys. Rev. Lett. **86**, 770 (2001), hep-ex/0007028.
 - [3] M. Ablikim et al. (BES) (2005), hep-ex/0506055.
 - [4] L. Maiani, F. Piccinini, A. D. Polosa, and V. Riquer, Phys. Rev. Lett. **93**, 212002 (2004), hep-ph/0407017.
 - [5] N. A. Tornqvist, Z. Phys. **C68**, 647 (1995), hep-ph/9504372.
 - [6] R. L. Jaffe, Phys. Rev. **D15**, 267 (1977).
 - [7] R. L. Jaffe, Phys. Rev. **D15**, 281 (1977).
 - [8] F. E. Close and N. A. Tornqvist, J. Phys. **G28**, R249 (2002), hep-ph/0204205.
 - [9] J. D. Weinstein and N. Isgur, Phys. Rev. Lett. **48**, 659 (1982).
 - [10] J. D. Weinstein and N. Isgur, Phys. Rev. **D27**, 588 (1983).
 - [11] J. D. Weinstein and N. Isgur, Phys. Rev. **D41**, 2236 (1990).
 - [12] R. L. Jaffe and F. E. Low, Phys. Rev. **D19**, 2105 (1979).
 - [13] M. G. Alford and R. L. Jaffe, Nucl. Phys. **B578**, 367 (2000), hep-lat/0001023.
 - [14] H. J. Lipkin, Phys. Lett. **B70**, 113 (1977).
 - [15] D. Black, A. H. Fariborz, F. Sannino, and J. Schechter, Phys. Rev. **D59**, 074026 (1999), hep-ph/9808415.
 - [16] J. R. Pelaez, Phys. Rev. Lett. **92**, 102001 (2004), hep-ph/0309292.
 - [17] D. M. Brink and F. Stancu, Phys. Rev. **D57**, 6778 (1998).
 - [18] D. M. Brink and F. Stancu, Phys. Rev. **D49**, 4665 (1994).
 - [19] H.-M. Chan and H. Hogaasen, Phys. Lett. **B72**, 121 (1977).
 - [20] R. L. Jaffe, Phys. Rept. **409**, 1 (2005), hep-ph/0409065.
 - [21] C. Amsler and N. A. Tornqvist, Phys. Rept. **389**, 61 (2004).
 - [22] S. Eidelman et al. (Particle Data Group), Phys. Lett. **B592**, 1 (2004), Particle Data Group collaboration.
 - [23] F. Iachello, N. C. Mukhopadhyay, and L. Zhang, Phys. Rev. **D44**, 898 (1991).
 - [24] F. J. Llanes-Estrada, ECONF **c0309101**, FRWP011 (2003), hep-ph/0311235.
 - [25] R. L. Jaffe, Phys. Rev. **D17**, 1444 (1978).
 - [26] F. Iachello, N. C. Mukhopadhyay, and L. Zhang, Phys. Lett. **B256**, 295 (1991).

- [27] F. Gursey and L. A. Radicati, Phys. Rev. Lett. **13**, 173 (1964).
- [28] F. Gursey, T. D. Lee, and M. Nauenberg, Phys. Rev. **135**, B467 (1964).
- [29] A. De Rujula, H. Georgi, and S. L. Glashow, Phys. Rev. **D12**, 147 (1975).
- [30] S. Okubo, Phys. Lett. **5**, 165 (1963).
- [31] G. Zweig (1964), CERN-TH-412.
- [32] J. Iizuka, K. Okada, and O. Shito, Prog. Theor. Phys. **35**, 1061 (1966).
- [33] J. J. Hernandez et al., Phys. Lett. **B239**, 1 (1990).
- [34] T. Regge, Nuovo Cim. **14**, 951 (1959).
- [35] K. Johnson and C. B. Thorn, Phys. Rev. **D13**, 1934 (1976).
- [36] G. 't Hooft, Nucl. Phys. **B75**, 461 (1974).
- [37] S. Godfrey and N. Isgur, Phys. Rev. **D32**, 189 (1985).
- [38] E. M. Aitala et al. (E791), Phys. Rev. Lett. **89**, 121801 (2002), hep-ex/0204018.
- [39] M. Neubert and B. Stech, Phys. Lett. **B231**, 477 (1989).
- [40] M. Neubert and B. Stech, Phys. Rev. **D44**, 775 (1991).
- [41] F. E. Close and A. W. Thomas, Phys. Lett. **B212**, 227 (1988).
- [42] M. G. Alford, K. Rajagopal, and F. Wilczek, Phys. Lett. **B422**, 247 (1998), hep-ph/9711395.
- [43] A. Selem and F. Wilczek (2006), hep-ph/0602128.
- [44] T. Schafer, Phys. Rev. **D68**, 114017 (2003), hep-ph/0309158.
- [45] P. Faccioli, Int. J. Mod. Phys. **A20**, 4615 (2005), hep-ph/0411088.
- [46] M. Oka (2004), hep-ph/0411320.
- [47] T. Schafer and E. V. Shuryak, Rev. Mod. Phys. **70**, 323 (1998), hep-ph/9610451.
- [48] C. Alexandrou, P. de Forcrand, and B. Lucini, PoS **LAT2005**, 053 (2006), hep-lat/0509113.
- [49] D. N. Walters, Nucl. Phys. Proc. Suppl. **119**, 553 (2003), hep-lat/0208011.
- [50] M. Hess, F. Karsch, E. Laermann, and I. Wetzorke, Phys. Rev. **D58**, 111502 (1998), hep-lat/9804023.
- [51] R. Alkofer, M. Kloker, A. Krassnigg, and R. F. Wagenbrunn, Phys. Rev. Lett **96**, 022001 (2006), hep-ph/0510028.
- [52] L. Y. Glozman and K. Varga, Phys. Rev. **D61**, 074008 (2000), hep-ph/9901439.
- [53] D. B. Leinweber, Phys. Rev. **D47**, 5096 (1993), hep-ph/9302266.
- [54] R. L. Jaffe (1999), hep-ph/0001123.
- [55] D. B. Lichtenberg, R. Roncaglia, and E. Predazzi (1996), hep-ph/9611428.
- [56] N. Mathur et al. (2006), hep-ph/0607110.
- [57] A. De Rujula, H. Georgi, and S. L. Glashow, Phys. Rev. Lett. **38**, 317 (1977).
- [58] A. Aloisio et al. (KLOE), Phys. Lett. **B536**, 209 (2002), hep-ex/0204012.
- [59] I. Caprini, G. Colangelo, and H. Leutwyler (2005), hep-ph/0512364.
- [60] D. V. Bugg (2005), hep-ex/0510021.
- [61] T. A. Kaeding (1995), nucl-th/9502037.
- [62] J. J. de Swart, Rev. Mod. Phys. **35**, 916 (1963).
- [63] This nonet has quantum numbers $n = L_{12-34} = S_{tot} = J = 0$, so we do not need to know the parameters a , b , c and d
- [64] This fit has been accomplished in a weighted way: for $a_0(980)$ and $f_0(980)$ the standard weights, coming from the inverse of the squared errors, have been used; the PDG does not give an estimate of the average values of the masses for $f_0(600)$ and $\kappa(800)$ and the values from different experiments are scattered in a range of $400 \div 700 \text{ MeV}$ and $700 \div 900 \text{ MeV}$ respectively, which have been used for the calculation of the weights, as an estimate of their unreliability.